

Review Article

Nutmeg-vertebrate interactions in the Asia-Pacific region: importance of frugivores for seed dispersal in Myristicaceae

Shumpei Kitamura¹ and Pilai Poonswad²

¹ Department of Environmental Science, Faculty of Bioresources and Environmental Sciences, Ishikawa Prefectural University, 1-308, Suematu, Nonoichi, Ishikawa, 921-8836, Japan

² Department of Microbiology, Faculty of Science, Mahidol University, Rama 6 Rd., Bangkok 10400, Thailand

Corresponding author: E-mail address: shumpei@ishikawa-pu.ac.jp

Abstract

In tropical forests, large frugivores are assumed to be important seed dispersers for many large-seeded trees such as the Myristicaceae, a widespread and common family. However, not all frugivores are effective seed dispersers, and understanding which frugivores are effective is vital for conservation biology. Here, we summarize the available data on fruit characteristics and frugivores for a large number of Myristicaceae species in the Asia-Pacific region and suggest future directions for evaluating the effects of disperser loss for these trees. Studies of fruit characteristics of Myristicaceae were highly biased toward morphological information, and few studies examined reproductive phenology or fruit chemistry. We identified 338 instances of nutmeg-frugivore interactions that included 129 species of Myristicaceae and 109 species of frugivores, including 40 bird, 68 mammal, and one reptile species. Large birds were major seed dispersers for this tree family. These bird species, such as hornbills and pigeons, consumed a variety of nutmeg species, remained briefly at fruiting trees, and dispersed intact seeds far from the parent trees in the forest. Although most seeds dispersed by birds subsequently suffered high seed predation by rodents, some germinated and established as seedlings, indicating the qualitative effectiveness of large birds as seed dispersers for Myristicaceae. Mammals were also major consumers of Myristicaceae. Gibbons, macaques, and civets potentially acted as long-distance dispersers for some nutmeg species. Orangutans, leaf monkeys, squirrels, and rodents consumed a variety of nutmeg species, but their roles as seed dispersers for Myristicaceae remain unclear. Studies of nutmeg-vertebrate interactions have typically focused on frugivory, whereas few studies have specifically quantified the effectiveness of frugivores as seed dispersers; thus, it remains difficult to evaluate the effect of frugivore loss on the populations of most nutmeg species in this region. Further studies of nutmeg-frugivore interactions are of great ecological importance, and the results of such studies will contribute to a general understanding of which evolutionary forces may have shaped current nutmeg-frugivore interactions in tropical forests worldwide.

Key words: *Endocomia*, *Gymnacranthera*, *Horsfieldia*, *Knema*, *Myristica*, seed predation

Received: 19 September 2012; Accepted: 21 May 2013; Published: 11 November 2013.

Copyright: Shumpei Kitamura and Pilai Poonswad. This is an open access paper. We use the Creative Commons Attribution 3.0 license <http://creativecommons.org/licenses/by/3.0/> - The license permits any user to download, print out, extract, archive, and distribute the article, so long as appropriate credit is given to the authors and source of the work. The license ensures that the published article will be as widely available as possible and that the article can be included in any scientific archive. Open Access authors retain the copyrights of their papers. Open access is a property of individual works, not necessarily journals or publishers.

Cite this paper as: Kitamura, S. and Poonswad, P. 2013. Nutmeg-vertebrate interactions in the Asia-Pacific region: importance of frugivores for seed dispersal in Myristicaceae. *Tropical Conservation Science*. Vol. 6(5):608-636. Available online: www.tropicalconservationscience.org

Introduction

Seed dispersal plays a critical role in the maintenance and recovery of plant diversity [1, 2]. This is especially true for the dispersal of seeds in highly diverse tropical rain forests, which usually support a wide range of potential dispersers. Most tropical woody plants have fleshy fruits [3, 4], and most tropical vertebrates eat fruits [5, 6]. In general, fruit (or seed) size usually limits the number of frugivores that can disperse the seeds [7-9]. Small fruits and large, soft fruits with many small seeds are consumed by a wide spectrum of frugivores, whereas larger fruits with a single large seed are consumed by relatively few potential dispersers [10, 11]. The seed dispersal of large-seeded plant species is therefore primarily dependent on large-bodied frugivores that are susceptible to extinction as a result of deleterious direct and indirect anthropogenic activities [9, 11-13].

The tropical rain forest tree family Myristicaceae consists of 500 species in 20 genera with a pantropical distribution centered in Malesia [14, 15]. In ecological studies of tropical rain forest tree communities, the Myristicaceae, or nutmeg, frequently ranks among one of the most important tree families, based on species frequency as well as species diversity [16]. In lowland rain forests of Malaysia, 30-40 nutmeg species have been recorded at a given study site, e.g., Lambir Hills National Park, Sarawak [17] and Pasoh Forest Reserve, Peninsular Malaysia [18]. Based on tree inventory data from 28 lowland dipterocarp rain forest locations throughout Borneo, Myristicaceae was the sixth most abundant tree family, accounting for 4.3% of all trees, and *Knema* was the eighth most common tree genus, accounting for 2.2% of trees in lowland dipterocarp forests [19]. The Myristicaceae is therefore an ideal family for studying general patterns of tropical forest diversity, and additional data for the nutmeg family will contribute valuable information to the general understanding of tropical forests.

Myristicaceae has a relatively conserved fruit and seed morphology (Fig. 1); therefore, one might expect that similar suites of frugivores consume and disperse the seeds. In the Neotropics, numerous studies related to the seed dispersal of Myristicaceae have been conducted since the 1970s, and large frugivorous birds and primates have been documented as the primary seed dispersers for this family [20]. In the Asia-Pacific region where the six genera *Endocomia*, *Gymnacranthera*, *Horsfieldia*, *Knema*, *Myristica*, and *Paramyristica* are distributed (Fig. 2), the primary seed-dispersal agents of these fruits are large birds such as fruit pigeons, hornbills, and birds of paradise [21-23]; however, the available data on seed dispersal in Myristicaceae are still limited, especially in terms of the effectiveness of dispersers [24]. Several criteria have been used to determine the efficiency and effectiveness of vertebrate frugivores as dispersers [24]; these include the size and diversity of fruits ingested, high fruit consumption, short visitation times, long gut retention times with seeds undamaged after gut passage, fruits swallowed whole with few dropped below parent trees,

behavior and movements during and after feeding, and seed deposition at suitable sites for germination.



Fig. 1. Ripe fruits of Myristicaceae in Budo-Sungai Padi National Park, southern Thailand. a: *Knema globularia*, b: *Horsfieldia tomentosa*, c: *Myristica iners*.

Although human impacts on frugivores specifically affect trees with large seeds in this region [11, 25], information on nutmeg-vertebrate interactions remains limited. Making existing data accessible to a wider audience is the first step toward applying what is known about the effectiveness of nutmeg-eating frugivores as seed dispersers for Myristicaceae and filling in gaps in knowledge. Our objectives in this study were to summarize the available data for Myristicaceae in the Asia-Pacific region in terms of (i) fruit traits and (ii) each frugivore in terms of the quantity and quality of dispersal, as well as to provide future directions for studies evaluating the effects of disperser loss for this family of trees.




Fig. 2. Distribution of Myristicaceae in the Asia-Pacific region and the major study sites cited in this review. NP: National Park, WS: Wildlife Sanctuary.

Literature survey

We first searched for data on characteristics of nutmeg fruits using Flora Malesiana [15], which covers 335 nutmeg species distributed in Indonesia, Malaysia, Burnei Darussalam, Singapore, the Philippines, and Papua New Guinea. We supplemented this search with regional floras, including Flora of Thailand [26], Flora of China [27], Flora of the Darwin Region [28], Myristicaceae of Papua [29], and other relevant studies in India and Tonga, to cover the distribution limit of Myristicaceae in this region (Fig. 2). To obtain the mean fruit/seed sizes of each nutmeg species, we averaged the minimum and maximum sizes reported in these studies. In most cases, these values were measured from herbarium specimens; therefore, the sizes calculated from these studies were likely smaller than those for fresh fruits/seeds in the field.

We then compiled a database on nutmeg species known to be consumed and dispersed by frugivores from the Asia-Pacific region. On 20 May 2011, we searched the Web of Science to obtain publications including the keywords “nutmeg”, “Myristicaceae” and several frugivores that occur in this region, such as “hornbill,” “pigeon,” “civet,” “gibbon,” “leaf monkey,” “macaque,” and “rodent.” We also examined recent reviews on frugivory by elephants [30], hornbills [31], gibbons [32], bears [33], and orangutans [34]. We supplemented these searches with literature cited by relevant studies and searches of regional journals as well as books on primates [35, 36] and pigeons [37]. Whenever possible, we recorded body mass data for each frugivore species from various sources [36-43] and the individual sources obtained in the literature survey. Values for body mass were the average for males and females [44].

We attempted to survey all major references, but the compiled lists were not exhaustive. The literature varies in clarity and quality; we excluded general accounts that list diets without giving the original source of information found in field guides and newsletters, but we included data found in books, journals, conference proceedings, and theses. These combined approaches resulted in data from a total of 84 publications, including 13 from books/book sections, 62 from peer-reviewed journals, one from conference proceedings, and eight from theses. A high proportion of studies included data on frugivory by hornbills (23 publications), followed by leaf monkeys (22), gibbons (15), macaques (14), and pigeons (11). Several publications included frugivory data on different frugivore groups; thus, the total number of studies that focused on frugivory was over 84. Plant nomenclature was revised according to the International Plant Name Index.

Characteristics of fruits in Myristicaceae

Fruit and seed size

In total, 352 nutmeg species were compiled by literature survey, and data were obtained for fruit length and diameter (N = 322 spp.), seed length (134 spp.), and seed diameter (18 spp.). Fruit sizes of Myristicaceae strongly varied among genera and species. Mean fruit size (length × diameter) at the genus level occurred in the following order: *Gymnacranthera* (2.3 × 1.5 mm, 6 spp.), *Knema* (2.9 × 2.0 mm, 79 spp.), *Horsfieldia* (3.0 × 2.2 mm, 88 spp.), *Myristica* (4.5 × 2.9 mm, 142 spp.), *Paramyristica* (4.8 × 2.8 mm, 1 sp.), and *Endocomia* (5.2 × 2.5 mm, 4 spp.). Seed length of Myristicaceae also varied among genera and species (Fig. 3): *Endocomia* (2.4-4.5 mm, 4 spp.), *Horsfieldia* (0.9-5.5 mm, 4 spp.), *Knema* (2.1-2.5 mm, 4 spp.), *Paramyristica* (3.0 mm, 1 sp.), and *Myristica* (1.4-5.5 mm, 121 spp.).

The sizes of nutmeg fruits/seeds also varied among species within a study site. The mean diameters of flesh arillate seeds were 21 mm (14-29 mm, N = 24 spp.) in Kutai National Park, Borneo [45], and 20 mm (10-30 mm, N = 14 spp.) in Budo-Sungai Padi National Park, Thailand (S. Kitamura, unpublished data). In both study sites, the mean diameters of flesh arillate seeds of Myristicaceae

were significantly larger than those of non-Myristicaceae (Welch's t-test, $t = 5.70$, $P < 0.001$, nutmeg = 9 spp., non-nutmeg = 180 spp. in Kutai; $t = 2.97$, $P < 0.009$, nutmeg = 9 spp., non-nutmeg = 423 spp. in Budo-Sungai Padi). We did not find any study that focused on individual variation in fruit/seed sizes of Myristicaceae in this region.




Fig. 3. Seed size variation among genera in Myristicaceae.

Phenology

At the species level, most nutmeg species fruited every year in India [46-49], Thailand [50-52], the Philippines [53], and Australia [54]. In Khao Yai National Park, Thailand, *Knema elegans* fruited at the intraspecies level in all 6 years but varied among individuals [50]. At the same site, a different nutmeg species, *Horsfieldia amygdalina* (formerly reported as *Horsfieldia glabra* in [10]), fruited every year from 1996 to 2003, except for 2000 [52]. Similarly, some nutmeg species exhibited annual fruiting patterns in lowland dipterocarp forests of Borneo, but others had supra-annual fruiting patterns [55, 56]. However, most studies on the reproductive phenology of Myristicaceae from Bornean forests did not collect large enough sample sizes to measure individual variation within a species; thus, determining differences among fruiting behaviors of Myristicaceae in this region is still difficult.

Aril color

In the literature survey, aril color was reported for 61 nutmeg species (*Endocomia*: 3 spp., *Gymnacranthera*: 3 spp., *Horsfieldia*: 19 spp., *Knema*: 19 spp., and *Myristica*: 18 spp.). Of these, red was the most common color of nutmeg arils (45 spp.), followed by orange (26 spp.), yellow (9 spp.), and pink (2 spp.). Similarly, within a study site, red arils were the most common in Myristicaceae ($N = 7$ spp.), followed by orange (3 spp.) in Budo-Sungai Padi National Park, southern Thailand [57].

Fruit chemistry

Little research has been conducted on fruit chemistry of Myristicaceae in this region, but available data suggest that the seeds/arils of nutmegs contain high lipid content both in the seeds [58]: *Myristica elliptica* (55.0%), *Knema hookeriana* (28.4%), and *Myristica cinnamomea* (8.5%), and in the arils [21, 58, 59]: *M. elliptica* (15.8%), *Myristica* sp. (57%), *H. amygdalina* (41.6%), and *K. elegans* (18.2%). These high lipid contents were preferred by porcupines in one Malaysian forest [58].

Diversity of nutmeg species eaten by frugivores

Regarding the taxonomic representation of nutmeg species eaten by different frugivore groups, the numbers vary widely (Table 1). We identified 338 instances of nutmeg-frugivore interactions that included 129 species of Myristicaceae, including *Endocomia* (1 sp.), *Gymnacranthera* (5 spp.), *Horsfieldia* (15 spp.), *Knema* (31 spp.), *Myristica* (43 spp.), and unidentified Myristicaceae genera (33 spp.), as well as 109 species of frugivores, including 40 bird, 68 mammal, and 1 reptile species (Fig. 4, Appendix 1).




Fig. 4. The interactions between genera of Myristicaceae and frugivores in the Asia-Pacific region. Five genera and ‘unknown’ Myristicaceae sp. are eaten by 23 different frugivore groups (mammals and reptiles in grey, and birds in black). Width of triangles indicates the number of documented interactions.

In terms of the number of nutmeg species eaten by each frugivore group, we identified hornbills and pigeons as major consumers of Myristicaceae among birds, and primates and squirrels among mammals (Table 1). More complete diet information is available for more thoroughly studied frugivores. Furthermore, frugivores from less diverse ecosystems, such as dry forests of India, show lower nutmeg diversity in their diet. Only 27% of frugivores consumed three or more nutmeg species as part of their diets (Fig. 5a). The maximum number of nutmeg species eaten by one type of frugivore was achieved by the orangutan *Pongo pygmaeus* (27 species), followed by bushy-crested hornbills *Anorrhinus galeritus* (20), wreathed hornbills *Rhyticeros undulatus* (19), and great hornbills *Buceros bicornis* (15). We found no relationship between mean body mass and the number of nutmeg species eaten by any frugivore group (Spearman's rank correlation, $P > 0.05$).

The maximum number of frugivores that consumed specific nutmeg species was recorded for *K. elegans* (12 spp.), followed by *Horsfieldia irya* (11) and *M. elliptica* (11). Most nutmeg species (68.2%) were eaten by only one or two frugivore species (Fig. 5b). As expected, frugivores in forests with high nutmeg diversity or inhabiting long-term study sites consumed the largest numbers of nutmeg species (e.g., Borneo and southern Thailand; Fig. 2). However, the number of frugivore species recorded for a given nutmeg species was likely underestimated, as most data on nutmeg-frugivore interactions reported from this region are based on studies of frugivores and not on observations of fruiting nutmeg trees.

Table 1. Summary of nutmeg-frugivore interactions in Asia-Pacific region. Frugivore groups are listed in decreasing order of the number of nutmeg species eaten by each group.

Frugivore group	Body weight (kg)	No. of species	Genus					No. of nutmeg species	
			<i>Endocomia</i>	<i>Gymnacranthera</i>	<i>Horsfieldia</i>	<i>Knema</i>	<i>Myristica</i>		Unknown
Hornbill (Bucerotidae)	0.24-2.9	19	X	X	X	X	X	X	50
Pigeon (Columbidae)	0.07-0.6	11			X	X	X	X	21
Bird of Paradise (Paradisaeidae)	0.09-0.22	6					X		3
Myna (Sturnidae)	0.16	1			X		X		2
Broadbill (Eurylaimidae)	0.06	1				X			1
Junglefowl (Phasianidae)	0.58	1			X				1
Parrot (Psittacidae)	0.23	1					X		1
Bird Total		40	X	X	X	X	X	X	67
Leaf monkey (Cercopithecidae)	5.8-15.6	21			X	X	X	X	31
Orangutan (Hominidae)	57.5	1		X	X	X	X	X	24
Macaque (Cercopithecidae)	4.5-15	11			X	X	X	X	23
Squirrel (Sciuridae)	0.1-2	6			X	X	X		23
Gibbon (Hylobatidae)	5.7-12.4	9		X	X	X	X	X	15
Rodent (Muridae)	0.17-0.8	5			X	X	X		8
Porcupine (Hystricidae)	2.5-8	3			X	X	X		7
Bear (Ursidae)	46	1			X	X			3
Civet (Viverridae)	3-6.3	4				X	X		3
Mousedeer (Tragulidae)	2.3-7.5	2			X	X	X		3
Tupai (Tupaïidae)	0.17	1				X	X		2
Flying fox (Pteropodidae)	0.6	1					X		1
Deer (Cervidae)	24	1				X			1
Mongoose (Herpestidae)	2.7	1					X		1
Wild pig (Suidae)	70	1				X			1
Mammal Total		68		X	X	X	X	X	85
Lizard (Varanidae)	7	1			X		X		3
Reptile Total		1			X		X		3
Frugivore Total		109							129

Fruit tree visitation, seed retention time, and dispersal distance

Several studies observed fruit consumed from large-seeded trees in this region [13, 60, 61], but most did not include Myristicaceae; thus, the data available on the visit frequency of frugivores to nutmeg trees are limited. One of the best-observed nutmeg species was *Myristica hypargyrea* in Tonga [62]. These authors found that the Pacific pigeon *Ducula pacifica* (118 visits) visited most frequently, followed by the red shining musk parrot *Prosopiea tabuensis* (17 visits) and the insular flying fox *Pteropus tonganus* (5 visits). Most seeds of *M. hypargyrea* were estimated to be dispersed by *D. pacifica*; ca. 80% of those ingested were expelled directly beneath conspecific fruiting crowns, 20% were dispersed locally, and <0.3% were dispersed more than 300 m into a different forest type [62].

In a Bornean forest with diverse nutmeg species (12 species over 205 observation hours), Leighton found that the mean visitation rate of hornbills ranged from 0.02 to 0.72 visits per hour per crop, which was one of the most frequently visited fruiting trees by hornbills [45]. However, the author presented no data on visit lengths of hornbills or the number of arillate seeds eaten by hornbills per visit at nutmeg trees [45]; thus, we could not quantitatively estimate the effectiveness of hornbills from these data. Hornbills usually remained in fruiting trees for a median of 20 min [45, 47, 60, 61, 63], and the median seed retention time of *H. amygdalina* by captive hornbills (*B. bicornis* and *R. undulatus*) was 40-50 min; 22% of the seeds were retained longer than 1 h (S. Kitamura, unpublished data). As hourly movements of hornbills sometimes exceed 6 km in the wild (S. Kitamura, unpublished data), they are able to transport nutmeg seeds far from parent trees.

Other frugivores such as gibbons, macaques, and civets are also likely to disperse seeds 100 m beyond parent plants [64-68]; thus, they potentially act as long-distance dispersers for some nutmeg species. Squirrels sometimes carry a whole fruit in their mouth away from the Myristicaceae tree for consumption in the canopy of neighboring trees or for later consumption (M. Yasuda, personal communication), but they appear to be poor distance-dispersers, with seeds deposited no more than 10 m from the fruiting crown [69].




Fig. 5a Left. Frequency histogram of cumulative number of frugivores per nutmeg species in the Asia-Pacific region.
Fig. 5b Right. Frequency histogram of cumulative number of nutmeg species per frugivore species in the Asia-Pacific region.

Extent of damage to seeds during consumption

We found that hornbills and pigeons as well as gibbons, macaques, and civets were reported to disperse intact seeds of Myristicaceae [10, 46, 70-72]. Hornbills and pigeons (*Ducula* spp. and *Ptilinopus* spp.) mostly swallowed arillate seeds of Myristicaceae and regurgitated the seeds intact [10, 21, 62, 70]. Of the mammals, gibbons (*Hylobates lar*) and civets (*Viverra zibetha*) in Thailand swallow both the seed and aril of *K. elegans* and then disperse the seeds through defecation [71]. Long-tailed macaques (*Macaca fascicularis*) in Singapore spit out seeds of *Knema latericia* and *Knema laurina* [72]. Other primates, such as orangutans and leaf monkeys, squirrels, and rodents consume many Myristicaceae fruits (Table 1), but the literature often does not make a clear distinction between consumption of the pulp and seeds; thus, their roles as seed dispersers for Myristicaceae remain unclear.

Despite the long list of frugivory records (Appendix 1), most studies did not investigate the potential viability of the nutmeg seeds defecated, regurgitated, spat, or dropped away from the parent trees by frugivores. The available data do not suggest any negative effects on the germination success of nutmeg seeds regurgitated by hornbills. The germination success of *Horsfieldia kingii* was similar between hornbill-regurgitated seeds (41%) and fallen seeds (33%) in India [47]. Over 90% germination success was recorded for hornbill-regurgitated nutmeg seeds, including *Horsfieldia tomentosa*, *Knema globularia*, and *Myristica iners*, but less than 10% success for *H. amygdalina* in Thailand (S. Kitamura, unpublished data). Under laboratory conditions, the germination success of nutmeg seeds was relatively high [73]: *Gymnacranthera eugeniifolia* (88%, N = 100), *Knema curtisii* (90%, N = 20), *K. laurina* (98%, N = 150), *Myristica crassa* (44%, N = 16), *Myristica malaccensis* (40%, N = 5), and *Myristica malaccensis* (73%, N = 55), except for *Knema scortechinii* (8%, N = 50). Removal of the aril before planting was advantageous for *K. laurina* (98% germination success without arils vs. 20% with arils), but not for *Knema furfuracea* (100% germination success, N = 50 seeds with arils); thus, the effects of aril removal by frugivores before seed deposition may vary among nutmeg species.

Fate of dispersed seeds

Frugivore activity is difficult to follow in the field, especially for canopy-dwelling animals. One method is to compare the survival and germination of experimentally placed seeds to mimic seed dispersal by frugivores, and another is to follow the fates of seeds dispersed by frugivores in a particular area, such as nest trees of hornbills. In a seed-removal experiment for *M. hypargyrea* on the ground in Tonga [62], the authors found that most seeds had been removed or killed by rats. Most *Myristica* seeds predicted to establish (6.5%) were dispersed by *D. pacifica*, which is more than twice the percentage of undispersed seeds that established (2.9%) around fruiting trees [62].

Hornbill-dispersed seeds around hornbill nests and roost trees were not particularly suitable for seed establishment and recruitment in general [47, 74-76]. In India, for example, hornbill-dispersed seeds of *H. kingii* around nest trees mostly died due to high seed predation by porcupines [77]. At the same site, *H. kingii* did not recruit near parent trees and showed increased seed mortality by seed-eating mammals with increasing seed density [78]. Hornbill-dispersed seeds of *H. amygdalina* were often consumed before germination at hornbill nest and roost trees in Thailand [74, 75], but some seedlings survived over 32 months (S. Kitamura, unpublished data). In Sulawesi, seedlings of *Horsfieldia brachiata* were one of the five most common seedling species around the nesting trees of the red-knobbed hornbill *Rhyticeros cassidix* [76]. Most nutmeg seeds with high seed density are likely to be eaten and consequently die before germination, but some seeds scattered by birds are likely to germinate and survive at their deposited sites, indicating their qualitative effectiveness as seed dispersers for Myristicaceae.

Summary and future directions

Many studies have reported nutmeg consumption by various kinds of frugivores in the Asia-Pacific region. Currently available data suggest that large, canopy-dwelling birds, including hornbills and pigeons, may function as major seed-dispersal agents for the Myristicaceae. Of these, the potential of hornbills as seed dispersers for large-seeded plants is well documented [31, 79]. Moreover, these birds originated in the mid-Eocene and might have assisted in the rapid colonization of large-seeded plants, including the Myristicaceae [80]. Some nutmeg seeds dispersed by hornbills germinated and survived as seedlings for several years. Based on the results summarized here, hornbills provide excellent seed-dispersal services to the Myristicaceae in this region.

Hornbills are extensively hunted by humans for their beaks, feathers, casque, and meat, and the densities and species richness of many hornbill species have now been reduced at many sites in this region [79, 81-85]. We expect that in areas in which the diversity of frugivores, particularly large avian frugivores like hornbills, has been reduced, macaques, civets, and squirrels may be the only remaining seed-dispersal agents for nutmeg trees. Indeed, some large-seeded *Myristica* trees are showing evidence of the lack of dispersal agents in Singapore [12, 86] and Lambir Hills National Park, Sarawak, Malaysia (K. Kimura, personal communication); hornbills have vanished or are nearly extinct at these sites, and other large frugivores, such as imperial pigeons (*Ducula* spp.), are very rare. In a Malaysian forest in Pasoh where at least six species of hornbills were originally distributed [87], only pig-tailed macaques (*Macaca nemestrina*) and several species of *Callosciurus* squirrels visited nutmeg trees in the canopy (M. Yasuda, personal communication). Can nutmeg trees still be dispersed by these remaining frugivores? We could not find detailed data on fruit removal for most nutmeg species in this region; thus, this question is still difficult to answer. Recent studies of other tree species in this region have revealed that frugivores that service the same plant may differ greatly in seed-dispersal effectiveness [88, 89]; therefore, similar results are expected in Myristicaceae.

Current studies on nutmeg-frugivore interactions typically focus on frugivory by certain groups of frugivores, whereas few studies quantify their effectiveness as seed dispersers for Myristicaceae. Thus, evaluating the effect of dispersal loss for this tree family in this region remains challenging. We suggest the following recommendations for future research on seed dispersal of these trees in this region. First, more information must be collected on the natural history of the dispersal ecology of the Myristicaceae. The most obvious natural history gap concerns fruit removal by each frugivore group at fruiting trees. Without such data, the effectiveness of nutmeg consumers as seed dispersers cannot be quantified. Second, many frugivores consume fallen fruits on the ground [6, 58, 90], and some potentially act as secondary dispersers for large-seeded plants [61, 91-93]. Although available data suggest that nutmeg seeds are often eaten by rodents on the ground [62, 77, 78], understanding the roles of these secondary dispersers that are tolerant of anthropogenic changes is becoming increasingly important. Third, comparative studies are needed of the dispersal ecology of sympatric nutmeg species with different seed sizes at a range of sites representing major forest types and a variety of human impacts. Nutmeg species with small seeds are expected to be dispersed by a broad assemblage of frugivores, and declines in seed removal by large frugivores might be compensated for by increases in seed removal by small frugivores. In addition, genetic tools can also be used to determine the origin/sources (maternal trees) for established recruitment and trees to determine the seed shadows generated by past and current frugivores.

Nutmeg trees are relatively abundant and well described in the literature, and most species are easily recognized in the field [16]. As elucidated by this review, further studies of nutmeg-frugivore interactions will be of great ecological importance in this region, and the results of future studies will contribute to a general understanding of which evolutionary forces might have shaped current nutmeg-frugivore interactions in tropical forests worldwide.

Acknowledgements

We thank anonymous reviewers, Pierre-Michel Forget, and Simon A. Queenborough for their comments on the manuscript. This study was supported by the Mahidol University Government Research Grant, the National Center for Genetic Engineering and Biotechnology (BIOTEC), the Hornbill Research Foundation, and the JSPS Research Fellowships for S.K.

References

- [1] Howe, H. F. and Smallwood, J. 1982. Ecology of seed dispersal. *Annual Review of Ecology and Systematics* 13:201-228.
- [2] Herrera, C. M., ed. 2002 *Seed dispersal by vertebrates*. Blackwell Science, Massachusetts.
- [3] Turner, I. M. 2001. *The ecology of trees in the tropical rain forest*. Cambridge University Press, Cambridge.
- [4] Jordano, P. 2000. Fruits and frugivory. Pp. 125-166 in Fenner, M. (ed.) *Seeds: The Ecology of Regeneration in Plant Communities*. CABI Publishing, CAB International, Wallingford, UK.
- [5] Fleming, T. H., Breitwisch, R. and Whitesides, G. H. 1987. Patterns of tropical vertebrate frugivore diversity. *Annual Review of Ecology and Systematics* 18:91-109.
- [6] Corlett, R. T. 1998. Frugivory and seed dispersal by vertebrates in the Oriental (Indomalayan) Region. *Biological Reviews* 73:413-448.
- [7] Wheelwright, N. T. 1985. Fruit size, gape width, and the diets of fruit-eating birds. *Ecology* 66:808-818.
- [8] McConkey, K. R. and Drake, D. R. 2002. Extinct pigeons and declining bat populations: are large seeds still being dispersed in the tropical Pacific? Pp. 381-395 in Levey, D. J., Silva, W. R. and Galetti, M. (eds.) *Seed dispersal and frugivory: ecology, evolution and conservation*. CABI Publishing, New York.
- [9] Peres, C. A. and van Roosmalen, M. 2002. Primate frugivory in two species-rich Neotropical forests: implications for the demography of large-seeded plants in overhunted areas. Pp. 407-421 in Levey, D. J., Silva, W. R. and Galetti, M. (eds.) *Seed dispersal and frugivory: ecology, evolution and conservation*. CABI Publishing, New York.
- [10] Kitamura, S., Yumoto, T., Poonswad, P., Chuailua, P., Plongmai, K., Maruhashi, T. and Noma, N. 2002. Interactions between fleshy fruits and frugivores in a tropical seasonal forest in Thailand. *Oecologia* 133:559-572.
- [11] Hamann, A. and Curio, E. 1999. Interactions among frugivores and fleshy fruit trees in a Philippine submontane rainforest. *Conservation Biology* 13:766-773.
- [12] Corlett, R. T. 2002. Frugivory and seed dispersal in degraded tropical East Asian landscapes. Pp. 451-465 in Levey, D. J., Silva, W. R. and Galetti, M. (eds.) *Seed dispersal and frugivory: ecology, evolution and conservation*. CABI Publishing, New York.
- [13] Sethi, P. and Howe, H. F. 2009. Recruitment of hornbill-dispersed trees in hunted and logged forests of the Indian Eastern Himalaya. *Conservation Biology* 23:710-718.
- [14] Whitmore, T. C., ed. 1972 *Tree Flora of Malaya*. Vol. 1. Longman, Kuala Lumpur and London.
- [15] De Wilde, W. J. J. O. 2000. *Myristicaceae*. National Herbarium of the Netherlands, Leiden.
- [16] LaFrankie, J. V. 2010. *Trees of Tropical Asia: an illustrated guide to diversity*. Black Tree Publications, Inc., Bacnotan, Philippines.

- [17] Lee, H. S., Davies, S. J., LaFrankie, J. V., Tan, S., Yamakura, T., Itoh, A., Ohkubo, T. and Ashton, P. S. 2002. Dipterocarp forest in Lambir Hills National Park, Sarawak, Malaysia. *Journal of Tropical Forest Science* 14:379-400.
- [18] Davies, S. J., Noor, N. S. M., LaFrankie, J. V. and Ashton, P. S. 2003. The Tree of Pasoh Forest: Stand Structure and Floristic Composition of the 50-ha Forest Research Plot. Pp. 35-50 in Okuda, T., Manokaran, N., Matsumoto, Y., Niiyama, K., Thomas, S. C. and Ashton, P. S. (eds.) *Pasoh: ecology of a lowland rain forest in Southeast Asia*. Springer-Verlag, Tokyo, Japan.
- [19] Slik, J. W. F., Poulsen, A. D., Ashton, P. S., Cannon, C. H., Eichhorn, K. A. O., Kartawinata, K., Lanniari, I., Nagamasu, H., Nakagawa, M., van Nieuwstadt, M. G. L., Payne, J., Purwaningsih, Saridan, A., Sidiyasa, K., Verburg, R. W., Webb, C. O. and Wilkie, P. 2003. A floristic analysis of the lowland dipterocarp forests of Borneo. *Journal of Biogeography* 30:1517-1531.
- [20] Howe, H. F. 1981. Dispersal of a Neotropical nutmeg (*Virola sebifera*) by birds. *Auk* 98:88-98.
- [21] Beehler, B. M. and Dumbacher, J. P. 1996. More examples of fruiting trees visited by predominantly by birds of paradise. *Emu* 96:81-88.
- [22] Sinclair, J. 1958. A revision of the Malayan Myristicaceae Gard. *Gardens' Bulletin Singapore* 16:205-466.
- [23] Ridley, H. N. 1930. *The dispersal of plants throughout the world*. L. Reeve & Co., Ashford.
- [24] Schupp, E. W. 1993. Quantity, quality and the effectiveness of seed dispersal by animals. *Vegetatio* 108:15-29.
- [25] Kitamura, S., Suzuki, S., Yumoto, T., Chuailua, P., Plongmai, K., Poonswad, P., Noma, N., Maruhashi, T. and Suckasam, C. 2005. A botanical inventory of a tropical seasonal forest in Khao Yai National Park, Thailand: implications for fruit-frugivore interactions. *Biodiversity and Conservation* 14:1241-1262.
- [26] De Wilde, W. J. J. O. 2002. Myristicaceae. Pp. 42 in *Flora of Thailand*. Forest Herbarium, Department of National Parks, Wildlife and Plant Conservation Bangkok, Thailand.
- [27] Bingtao, L. and Wilson, T. K. 2008. Myristicaceae. Pp. 96-101 in *Flora of China Vol. 7 (Menispermaceae through Capparaceae)*. Missouri Botanical Garden Press, St. Louis.
- [28] Mangion, C. P. 2011. Myristicaceae. Pp. 1-5 in Short, P. S. and Cowie, I. D. (eds.) *Flora of the Darwin Region*. Northern Territory Herbarium, Department of Natural Resources, Environment, the Arts and Sport, Palmerston, N.T.
- [29] De Wilde, W. J. J. O. 2006. Myristicaceae of Papua. Pp. 408-415 in Marshall, A. J. and Beehler, B. M. (eds.) *The ecology of Papua*. Periplus Editions, Singapore.
- [30] Campos-Arceiz, A. and Blake, S. 2011. Megagardeners of the forest-the role of elephants in seed dispersal. *Acta Oecologica* 37:542-553.
- [31] Kitamura, S. 2011. Frugivory and seed dispersal by hornbills (Bucerotidae) in tropical forests. *Acta Oecologica* 37:531-541.
- [32] Elder, A. A. 2009. Hylobatid diets revisited: The importance of body mass, fruit availability, and interspecific competition. Pp. 133-159 in Lappan, S. and Whittaker, D. J. (eds.) *The Gibbons: New perspectives on small ape socioecology and population biology*. Springer, New York.
- [33] Steinmetz, R. 2011. Ecology and distribution of sympatric Asiatic Black Bears and Sun Bears in the seasonally dry forests of Southeast Asia. Pp. 249-273 in McShea, W. J., Davies, S. J. and Bhumpakphan, N. (eds.) *The ecology and conservation of seasonally dry forests in Asia*. Washington Institute Scholarly Press, Washington, D.C., USA.
- [34] Russon, A. E., Wich, S. A., Ancrenaz, M., Kanamori, T., Knott, C. D., Kuze, N., Morrogh-Bernard, H. C., Pratje, P., Ramlee, H. and Rodman, P. 2009. Geographic variation in orangutan diets. Pp. 135-156 in Wich, S. A., Utami Atmoko, S. S., Mitra Setia, T. and van Schaik, C. P. (eds.) *Orangutans: Geographic variation in behavioral ecology and conservation*. Oxford University Press, New York.

- [35] Payne, J. B. 1980. Competitors. *Malayan Forest Primates: ten years' study in tropical rain forest*:261-277.
- [36] Campbell, C. J., Fuentes, A., MacKinnon, K. C., Panger, M. and Bearder, S. K., eds. 2007 *Primates in perspective*. Oxford University Press, Oxford, UK.
- [37] Gibbs, D., Barnes, E. and Cox, J. 2000. *Pigeons and doves: a guide to the pigeons and doves of the world*. Yale University Press, New Haven & London, UK.
- [38] Kemp, A. C. 1995. *The hornbills, Bucerotiformes*. Oxford University Press, Oxford.
- [39] Kemp, A. C. 2001. Family Bucerotidae (Hornbills). Pp. 436-520 in del Hoyo, J., Elliott, A. and Sargatal, J. (eds.) *Handbook of the Birds of the World. Vol. 6. Mousebirds to Hornbills*. Lynx Edicions, Barcelona.
- [40] Gupta, A. K. and Chivers, D. J. 1999. Biomass and use of resources in South and South-East Asian primate communities. Pp. 38-54 in Fleagle, J., Janson, C. and Reed, K. E. (eds.) *Primate communities*. Cambridge University Press, Cambridge.
- [41] Payne, J., Francis, C. M. and Phillips, K. 1985. *A field guide to the mammals of Borneo*. The Sabah Society, WWF Malaysia, Kota Kinabaru, Kuala Lumpur.
- [42] Wells, D. R. 1999. *The Birds of the Thai-Malay Peninsula: Vol. 1 Non-passerines*. Academic Press, London, UK.
- [43] Wells, D. R. 2007. *The Birds of the Thai-Malay Peninsula: Vol. 2 Passerines*. Christopher Helm, London, UK.
- [44] Forget, P. M., Dennis, A. J., Mazer, S. J., Jansen, P. A., Kitamura, S., Lambert, J. A. and Westcott, D. A. 2007. Seed allometry and disperser assemblages in tropical rain forests: a comparison of four floras on different continents. Pp. 5-36 in Dennis, A. J., Schupp, E. W., Green, R. J. and Westcott, D. A. (eds.) *Seed dispersal: Theory and its application in a changing world*. CAB International, Wallingford, UK.
- [45] Leighton, M. 1982. *Fruit resources and patterns of feeding, spacing and grouping among sympatric Bornean hornbills (Bucerotidae)*. PhD thesis, University of California, California.
- [46] Kannan, R. and James, D. A. 1999. Fruiting phenology and the conservation of the Great Pied Hornbill (*Buceros bicornis*) in the Western Ghats of southern India. *Biotropica* 31:167-177.
- [47] Datta, A. 2001. *An ecological study of sympatric hornbills and fruiting patterns in a tropical forest in Arunachal Pradesh*. PhD thesis, Saurashtra University, Rajkot.
- [48] Ganesh, T. and Davidar, P. 2005. Fruiting phenology and pre-dispersal seed predation in a rainforest in southern Western Ghats, India. *Tropical fruits and frugivores: the search for strong interactors*:139-154.
- [49] Sharma, M. V. and Shivanna, K. R. 2011. Pollinators, pollination efficiency and fruiting success in a wild nutmeg, *Myristica dactyloides*. *Journal of Tropical Ecology* 27:405-412.
- [50] Brockelman, W. Y. 2011. Rainfall patterns and unpredictable fruit production in seasonally dry evergreen forest and their effects on gibbons. Pp. 195-216 in McShea, W. J., Davies, S. J. and Bhumpakphan, N. (eds.) *The ecology and conservation of seasonally dry forests in Asia*. Washington Institute Scholarly Press, Washington, D.C., USA.
- [51] Suzuki, S. 2007. *Fruit utilization patterns of terrestrial frugivores and the population dynamics of small mammal community in a seasonal tropical forest in Khao Yai National Park, Thailand*. PhD Thesis, University of Shiga Prefecture, Hikone, Japan (in Japanese).
- [52] Plongmai, K., Poonswad, P., Sukkasem, C. and Chuailua, P. 2005. The availability of ripe fruits in the annual hornbill life cycle. Pp. 131-142 in Lum, S. and Poonswad, P. (eds.) *The ecology of hornbills: reproduction and populations*. Pimdee Karnpim Co. Ltd., Bangkok, Thailand.
- [53] Hamann, A. 2004. Flowering and fruiting phenology of a Philippine submontane rain forest: climatic factors as proximate and ultimate causes. *Journal of Ecology* 92:24-31.

- [54] Armstrong, J. E. and Irvine, A. K. 1989. Flowering, sex ratios, pollen-ovule ratios, fruit set, and reproductive effort of a dioecious tree, *Myristica insipida* (Myristicaceae), in two different rain forest communities. *American Journal of Botany* 76:74-85.
- [55] Brearley, F. Q., Proctor, J., Suriantata, Nagy, L., Dalrymple, G. and Voysey, B. C. 2007. Reproductive phenology over a 10-year period in a lowland evergreen rain forest of central Borneo. *Journal of Ecology* 95:828-839.
- [56] Sakai, S., Momose, K., Yumoto, T., Nagamitsu, T., Nagamasu, H., Hamid, A. A. and Nakashizuka, T. 1999. Plant reproductive phenology over four years including an episode of general flowering in a lowland dipterocarp forest, Sarawak, Malaysia. *American Journal of Botany* 86:1414-1436.
- [57] Kitamura, S., Thong-Aree, S., Madsri, S. and Poonswad, P. 2011. Characteristics of hornbill-dispersed fruits in dipterocarp forests of southern Thailand. *Raffles Bulletin of Zoology* S24:137-147.
- [58] Yasuda, M., Miura, S., Ishii, N., Okuda, T. and Hussein, N. A. 2005. Fallen fruits and terrestrial vertebrate frugivores: a case study in a lowland tropical rain forest in Peninsular Malaysia. Pp. 151-174 in Forget, P. M., Lambert, J. E., Hulme, P. E. and Vander Wall, S. B. (eds.) *Seed fate: predation, dispersal and seedling establishment*. CABI Publishing, Wallingford.
- [59] Kanwatanakid-Savini, C., Poonswad, P. and Savini, T. 2009. An assessment of food overlap between gibbons and hornbills. *Raffles Bulletin of Zoology* 57:189-198.
- [60] Kitamura, S., Suzuki, S., Yumoto, T., Poonswad, P., Chuailua, P., Plongmai, K., Maruhashi, T., Noma, N. and Suckasam, C. 2006. Dispersal of *Canarium euphyllum* (Burseraceae), a large-seeded tree species, in a moist evergreen forest in Thailand. *Journal of Tropical Ecology* 22:137-146.
- [61] Kitamura, S., Suzuki, S., Yumoto, T., Poonswad, P., Chuailua, P., Plongmai, K., Noma, N., Maruhashi, T. and Suckasam, C. 2004. Dispersal of *Aglaia spectabilis*, a large-seeded tree species in a moist evergreen forest in Thailand. *Journal of Tropical Ecology* 20:421-427.
- [62] Meehan, H. J., McConkey, K. R. and Drake, D. R. 2005. Early fate of *Myristica hypargyrea* seeds dispersed by *Ducula pacifica* in Tonga, Western Polynesia. *Austral Ecology* 30:374-382.
- [63] Suryadi, S., Kinnaird, M. F., O'Brien, T. G., Supriatna, J. and Somadikarta, S. 1994. Food preferences of the Sulawesi red-knobbed hornbill during the non-breeding season. *Tropical Biodiversity* 2:377-384.
- [64] McConkey, K. R. and Chivers, D. J. 2007. Influence of gibbon ranging patterns on seed dispersal distance and deposition site in a Bornean forest. *Journal of Tropical Ecology* 23:269-275.
- [65] Tsujino, R. and Yumoto, T. 2009. Topography-specific seed dispersal by Japanese macaques in a lowland forest on Yakushima Island, Japan. *Journal of Animal Ecology* 78:119-125.
- [66] Nakashima, Y. and Sukor, J. A. 2010. Importance of common palm civets (*Paradoxurus hermaphroditus*) as a long-distance disperser for large-seeded plants in degraded forests. *Tropics* 18:221-229.
- [67] Yumoto, T., Noma, N. and Maruhashi, T. 1998. Cheek-pouch dispersal of seeds by Japanese monkeys (*Macaca fuscata yakui*) on Yakushima Island, Japan. *Primates* 39:325-338.
- [68] Terakawa, M., Isagi, Y., Matsui, K. and Yumoto, T. 2009. Microsatellite analysis of the maternal origin of *Myrica rubra* seeds in the feces of Japanese macaques. *Ecological Research* 24:663-670.
- [69] Becker, P. and Wong, M. 1985. Seed dispersal, seed predation, and juvenile mortality of *Aglaia* sp. (Meliaceae) in lowland dipterocarp rainforest. *Biotropica* 17:230-237.
- [70] McConkey, K. R., Meehan, H. J. and Drake, D. R. 2004. Seed dispersal by Pacific Pigeons (*Ducula pacifica*) in Tonga, Western Polynesia. *Emu* 104:369-376.
- [71] Whittington, C. 1990. Seed dispersal by white-handed gibbons (*Hylobates lar*) in Khao Yai National Park, Thailand.

- [72] Lucas, P. W. and Corlett, R. T. 1998. Seed dispersal by long-tailed macaques. *American Journal of Primatology* 45:29-44.
- [73] Ng, F. S. P. 1980. Germination ecology of Malaysian woody plants. *Malaysian Forester* 43:406-437.
- [74] Kitamura, S., Yumoto, T., Noma, N., Chuailua, P., Maruhashi, T., Wohandee, P. and Poonswad, P. 2008. Aggregated seed dispersal by wreathed hornbills at a roost site in a moist evergreen forest of Thailand. *Ecological Research* 23:943-952.
- [75] Kitamura, S., Yumoto, T., Poonswad, P., Noma, N., Chuailua, P., Plongmai, K., Maruhashi, T. and Suckasam, C. 2004. Pattern and impact of hornbill seed dispersal at nest trees in a moist evergreen forest in Thailand. *Journal of Tropical Ecology* 20:545-553.
- [76] Kinnaird, M. F. 1998. Evidence for effective seed dispersal by the Sulawesi red-knobbed hornbill, *Aceros cassidix*. *Biotropica* 30:50-55.
- [77] Velho, N., Datta, A. and Isvaran, K. 2009. Effect of rodents on seed fate of five hornbill-dispersed tree species in a tropical forest in north-east India. *Journal of Tropical Ecology* 25:507-514.
- [78] Velho, N., Isvaran, K. and Datta, A. 2012. Rodent seed predation: effects on seed survival, recruitment, abundance, and dispersion of bird-dispersed tropical trees. *Oecologia*:in press.
- [79] Kinnaird, M. F. and O'Brien, T. G. 2007. *The ecology and conservation of Asian hornbills: farmers of the forest*. University Of Chicago Press, Chicago.
- [80] Viseshakul, N., Charoennitikul, W., Kitamura, S., Kemp, A., Thong-Aree, S., Surapunpitak, Y., Poonswad, P. and Ponglikitmongkol, M. 2011. A phylogeny of frugivorous hornbills linked to the evolution of Indian plants within Asian rainforests. *Journal of Evolutionary Biology* 24:1533-1545.
- [81] Pattanavibool, A. and Dearden, P. 2002. Fragmentation and wildlife in montane evergreen forests, northern Thailand. *Biological Conservation* 107:155-164.
- [82] Datta, A. 1998. Hornbill abundance in unlogged forest, selectively logged forest and a forest plantation in Arunchal Pradesh, India. *Oryx* 32:285-294.
- [83] Johns, A. D. 1987. The use of primary and selectively logged rain-forest by Malaysian hornbills (Bucerotidae) and implications for their conservation. *Biological Conservation* 40:179-190.
- [84] Johns, A. D. 1985. Selective logging and wildlife conservation in tropical rain-forest: problems and recommendations. *Biological Conservation* 31:355-375.
- [85] Gonzalez, J. C. T. 2011. Enumerating the ethno-ornithological importance of Philippine hornbills. *Raffles Bulletin of Zoology* S24:149-161.
- [86] Ercelawn, A. C., La Frankie, I. V., Lum, S. K. Y. and Lee, S. K. 1998. Short-term recruitment of trees in a forest fragment in Singapore. *Tropics* 8:105-115.
- [87] Francis, C. M. and Wells, D. R. 2003. The Birds Community at Pasoh: Composition and Population Dynamics. Pp. 375-393 in Okuda, T., Manokaran, N., Matsumoto, Y., Niiyama, K., Thomas, S. C. and Ashton, P. S. (eds.) *Pasoh: ecology of a lowland rain forest in Southeast Asia*. Springer-Verlag, Tokyo, Japan.
- [88] Brodie, J. F., Helmy, O. E., Brockelman, W. Y. and Maron, J. L. 2009. Functional differences within a guild of tropical mammalian frugivores. *Ecology* 90:688-698.
- [89] McConkey, K. R. and Brockelman, W. Y. 2011. Nonredundancy in the dispersal network of a generalist tropical forest tree. *Ecology* 92:1492-1502.
- [90] Prasad, S., Pittet, A. and Sukumar, R. 2010. Who really ate the fruit? A novel approach to camera trapping for quantifying frugivory by ruminants. *Ecological Research* 25:225-231.
- [91] Yasuda, M., Miura, M. and Hussein, N. A. 2000. Evidence for food hoarding behaviour in terrestrial rodents in Pasoh forest reserve, a Malaysian lowland rain forest. *Journal of Tropical Forest Science* 12:164-173.
- [92] Kitamura, S., Yumoto, T., Poonswad, P., Suzuki, S. and Wohandee, P. 2008. Rare seed-predating mammals determine seed fate of *Canarium euphyllum*, a large-seeded tree species in a moist evergreen forest, Thailand. *Ecological Research* 23:169-177.

- [93] Cao, L., Xiao, Z., Guo, C. and Chen, J. 2011. Scatter-hoarding rodents as secondary seed dispersers of a frugivore-dispersed tree *Scleropyrum wallichianum* in a defaunated Xishuangbanna tropical forest, China. *Integrative Zoology* 6:227-234.
- [94] Wich, S. A., Atmoko, S. S. U. and Setia, T. M., eds. 2009 *Orangutans: geographic variation in behavioral ecology and conservation*. Oxford University Press, USA.
- [95] Whitten, A. J. 1982. Diet and feeding-behavior of kloss gibbons on Siberut island, Indonesia. *Folia Primatologica* 37:177-208.
- [96] Kinnaird, M. F. and O'Brien, T. G. 1993. Preliminary observation on the breeding biology of the endemic Sulawesi red-knobbed hornbill (*Rhyticeros cassidix*). *Tropical Biodiversity* 1:107-112.
- [97] Chimchome, V., Vidhidharm, A., Simchareon, S., Bumrungsi, S. and Poonswad, P. 1998. Comparative study of the breeding biology and ecology of two endangered hornbill species in Huai Kha Khaeng wildlife sanctuary, Thailand. *The Asian hornbills: ecology and conservation, Thai Studies in Biodiversity No 2*:111-131.
- [98] Kitamura, S., Yumoto, T., Poonswad, P., Chuailua, P. and Plongmai, K. 2004. Characteristics of hornbill-dispersed fruits in a tropical seasonal forest in Thailand. *Bird Conservation International* 14:S81-S88.
- [99] Hadiprakarsa, Y. Y. and Kinnaird, M. F. 2004. Foraging characteristics of an assemblage of four Sumatran hornbill species. *Bird Conservation International* 14:S53-S62.
- [100] Harrison, M. E. 2009. *Orangutan feeding behaviour in Sabangau, Central Kalimantan*. PhD thesis, University of Cambridge, Cambridge.
- [101] Supriatna, J., Manullang, B. O. and Soekara, E. 1986. Group composition, home range, and diet of the maroon leaf monkey (*Presbytis rubicunda*) at Tanjung Puting Reserve, Central Kalimantan, Indonesia. *Primates* 27:185-190.
- [102] Poirier, F. E. and Smith, E. O. 1974. The Crab-Eating Macaques (*Macaca fascicularis*) of Angaur Island, Palau, Micronesia. *Folia primatol* 22:258-306.
- [103] Chivers, D. J., ed. 1980 *Malayan forest primates: ten years' study in tropical rain forest*. Plenum Press, New York, USA.
- [104] Gittins, S. P. 1982. Feeding and ranging in the agile gibbon. *Folia Primatologica* 38:39-71.
- [105] Jayasekara, P., Weerasinghe, U. R., Wijesundara, S. and Takatsuki, S. 2007. Identifying diurnal and nocturnal frugivores in the terrestrial and arboreal layers of a tropical rain forest in Sri Lanka. *Ecotropica* 13:7-15.
- [106] Marsden, S. J. and Pilgrim, J. D. 2003. Factors influencing the abundance of parrots and hornbills in pristine and disturbed forests on New Britain, PNG. *Ibis* 145:45-53.
- [107] Datta, A. and Rawat, G. S. 2003. Foraging patterns of sympatric hornbills during the nonbreeding season in Arunachal Pradesh, northeast India. *Biotropica* 35:208-218.
- [108] Datta, A. and Rawat, G. S. 2008. Dispersal modes and spatial patterns of tree species in a tropical forest in Arunachal Pradesh, northeast India. *Tropical Conservation Science* 1:163-185.
- [109] Solanki, G. S., Kumar, A. and Sharma, B. K. 2008. Feeding ecology of *Trachypithecus pileatus* in India. *International Journal of Primatology* 29:173-182.
- [110] Riley, E. P. 2007. Flexibility in diet and activity patterns of *Macaca tonkeana* in response to anthropogenic habitat alteration. *International Journal of Primatology* 28:107-133.
- [111] Fredriksson, G. M., Wich, S. A. and Trisno 2006. Frugivory in sun bears (*Helarctos malayanus*) is linked to El Nino related fluctuations in fruiting phenology, East Kalimantan, Indonesia. *Biological Journal of the Linnean Society* 89:489-508.
- [112] Mudappa, D. 2000. Breeding biology of the Malabar Grey Hornbill (*Ocyrceros griseus*) in southern Western Ghats, India. *Journal of Bombay Natural History Society* 97:15-24.
- [113] Mudappa, D., Kumar, A. and Chellam, R. 2010. Diet and fruit choice of the brown palm civet *Paradoxurus jerdoni*, a viverrid endemic to the Western Ghats rainforest, India. *Tropical Conservation Science* 3:282-300.

- [114] Punekar, S. A. 2002. Some food plants of Hanuman langur *Semnopithecus entellus* (Dufresne) in the western Ghats of Maharashtra, India. *ZOOS'PRINT JOURNAL* 17:797-801.
- [115] Umapathy, G. and Kumar, A. 2000. Impacts of the habitat fragmentation on time budget and feeding ecology lion-tailed macaque (*Macaca silenus*) in rain forest fragments of Anamalai Hills, South India. *Primate report* 58:67-82.
- [116] Farida, W. R. and Harun 2000. The diversity of plants and feed resources for the Java Gibbon (*Hylobates moloch*), Grizzled Langur (*Presbytis comata*), and Silver Langur (*Trachypithecus auratus*) in Gunung Halimun National Park. *Jurnal of Primatologi Indonesia* 3:55-61.
- [117] Bartlett, T. Q. 1999. Feeding and ranging behavior of the white-handed gibbon (*Hylobates lar*) in Khao Yai National Park, Thailand.
- [118] Srikosamatar, S. 1984. Ecology of pileated gibbons in south-east Thailand. Pp. 242-257 in Preuschoft, H., Chivers, D. J., Brockelman, W. Y. and Creel, N. (eds.) *The Lesser Apes: Evolutionary and Behavioural Biology*. Edinburgh.
- [119] Whittington, C. and Treesucon, U. 1991. Selection and treatment of food plants by white-handed gibbons (*Hylobates lar*) in Khao Yai National Park, Thailand. *Natural History Bulletin of the Siam Society* 39:111-122.
- [120] Leighton, M. 1993. Modeling dietary selectivity by Bornean orangutans: Evidence for integration of multiple criteria in fruit selection. *International Journal of Primatology* 14:257-313.
- [121] Lucas, P. W. and Corlett, R. T. 1991. Relationship between the diet of *Macaca fascicularis* and forest phenology. *Folia Primatologica* 57:201-215.
- [122] McConkey, K. R., Aldy, F., Ario, A. and Chivers, D. J. 2002. Selection of fruit by gibbons (*Hylobates muelleri* x *agilis*) in the rain forests of Central Borneo. *International Journal of Primatology* 23:123-145.
- [123] Kanamori, T., Kuze, N., Bernard, H., Malim, T. P. and Kohshima, S. 2010. Feeding ecology of Bornean orangutans (*Pongo pygmaeus morio*) in Danum Valley, Sabah, Malaysia: a 3-year record including two mast fruitings. *American Journal of Primatology* 72:820-840.
- [124] Ouithavon, K., Poonswad, P., Bhumbhakpan, N. and Laohajinda, V. 2005. A comparative study of the feeding ecology of two sympatric Hornbill species (Aves: Bucerotidae) during their breeding season in Huai Kha Khaeng Wildlife Sanctuary, Thailand. Pp. 59-73 in Lum, S. and Poonswad, P. (eds.) *The ecology of hornbills: reproduction and populations*. Pimdee Karnpim Co. Ltd., Bangkok, Thailand.
- [125] Ouithavon, K., Poonswad, P., Bhumbhakpan, N. and Laohajinda, V. 2005. Some characteristics of food of two sympatric hornbill species (Aves: Bucerotidae) and fruit availability during their breeding season in Huai Kha Khaeng Wildlife Sanctuary, Thailand. Pp. 75-85 in Lum, S. and Poonswad, P. (eds.) *The ecology of hornbills: reproduction and populations*. Pimdee Karnpim Co. Ltd., Bangkok, Thailand.
- [126] Ungar, P. S. 1995. Fruit preferences of four sympatric primate species at Ketambe, Northern Sumatra, Indonesia. *International Journal of Primatology* 16:221-245.
- [127] Ali, S. and Ripley, S. D. 1969. *Handbook of the birds of India and Pakistan. Vol 3*. Oxford University Press, London, UK.
- [128] Kauth, M., Engel, S., Lastimoza, L. L. and Curio, E. 1998. Observations on the breeding biology of the Writhed-billed Hornbill (*Aceros waldeni*) in the Philippines. *Journal Fur Ornithologie* 139:475-483.
- [129] Ganesh, T. 2004. Seed predation and seed removal rates in *Myristica dactylodes* in southern India. *Workshop on seed dispersal & frugivory in Asia*:11.
- [130] Umapathy, G. 1998. *Impacts of habitat fragmentation on the arboreal mammals in the wet evergreen forests of the Anamalai Hills in the Western Ghats, south India*. PhD thesis, Bharathiar University, Coimbatore.

- [131] Mack, A. L. and Wright, D. D. 1996. Notes on occurrence and feeding of birds at Crater Mountain Biological Research Station, Papua New Guinea. *Emu* 96:89-101.
- [132] Collar, N. J., Crosby, R. and Crosby, M. J., eds. 2001 *Threatened birds of Asia: the BirdLife International red data book*. BirdLife International, Cambridge, UK.
- [133] McConkey, K. R., Drake, D. R., Meehan, H. J. and Parsons, N. 2003. Husking stations provide evidence of seed predation by introduced rodents in Tongan rain forests. *Biological Conservation* 109:221-225.
- [134] Crome, F. H. J. 1975. The ecology of fruit pigeons in tropical northern Queensland. *Australian Wildlife Research* 2:155-185.
- [135] Ali, S. and Ripley, S. D. 1970. *Handbook of the birds of India and Pakistan. Vol 4*. Oxford University Press, London, UK.
- [136] Beehler, B. 1983. Frugivory and polygamy in birds of paradise. *Auk* 100:1-11.
- [137] Palombit, R. A. 1997. Inter- and intraspecific variation in the diets of sympatric siamang (*Hylobates syndactylus*) and lar gibbons (*Hylobates lar*). *Folia Primatologica* 68:321-337.
- [138] Smythies, B. E. 1960. *The birds of Borneo*. Oliver and Boyd, Edinburgh, UK.
- [139] Aziz, M. A. and Feeroz, M. M. 2009. Utilization of forest flora by Phayre's Leaf-Monkey *Trachypithecus phayrei* (Primates: Cercopithecidae) in semi-evergreen forests of Bangladesh. *Journal of Threatened Taxa* 1:257-262.
- [140] Bleisch, W. V. and Chen, N. 1991. Ecology and behavior of wild black-crested gibbons (*Hylobates concolor*) in China with a reconsideration of evidence for polygyny. *Primates* 32:539-548.
- [141] Dela, J. 2007. Seasonal food use strategies of a colobine frugivore, *Trachypithecus vetulus nestor*, at Panadura and Piliyandala, Sri Lanka. *International Journal of Primatology* 28:607-626.
- [142] Duc, H. M., Baxter, G. S. and Page, M. J. 2009. Diet of *Pygathrix nigripes* in southern Vietnam. *International Journal of Primatology* 30:15-28.
- [143] Fan, P., Ni, Q., Sun, G., Huang, B. and Jiang, X. 2009. Gibbons under seasonal stress: the diet of the black crested gibbon (*Nomascus concolor*) on Mt. Wuliang, Central Yunnan, China. *Primates* 50:37-44.
- [144] Frith, H. J., Crome, F. H. J. and Wolfe, T. O. 1976. Food of fruit-pigeons in New Guinea. *Emu* 76:49-58.
- [145] Fuentes, A. 1996. Feeding and ranging in the Mentawai Island langur (*Presbytis potenziani*). *International Journal of Primatology* 17:525-548.
- [146] Goodwin, D. 1977. *Pigeons and doves of the world*. British Museum (Natural History), London, UK.
- [147] Grueter, C. C., Da-Yong, L. I., Shun-Kai, F. and Bao-Ping, R. E. N. 2010. Niche partitioning between sympatric rhesus macaques and Yunnan snub-nosed monkeys at Baimaxueshan Nature Reserve, China. *Zoological Research* 5:516-522.
- [148] Gurmaya, K. J. 1986. Ecology and behavior of *Presbytis thomasi* in Northern Sumatra. *Primates* 27:151-172.
- [149] Islam, M. A. and Feeroz, M. M. 1992. Ecology of hoolock gibbon of Bangladesh. *Primates* 33:451-464.
- [150] Kool, K. M. 1993. The diet and feeding behavior of the silver leaf monkey (*Trachypithecus auratus sondaicus*) in Indonesia. *International Journal of Primatology* 14:667-700.
- [151] Krishnamani, R. 1994. Diet composition of the bonnet macaque (*Macaca radiata*) in a tropical dry evergreen forest of southern India. *Tropical Biodiversity* 2:285.
- [152] Kumar, A. and Solanki, G. S. 2008. Population status and conservation of capped langurs (*Trachypithecus pileatus*) in and around Pakke Wildlife Sanctuary, Arunachal Pradesh, India. *Primate Conservation* 23:97-105.

- [153] Kumar, R. S., Mishra, C. and Sinha, A. 2007. Foraging ecology and time-activity budget of the Arunachal macaque *Macaca munzala*: A preliminary study. *Current Science* 93:532-539.
- [154] Le Khac Quyet, N. A. D., Tai, V. A., Wright, B. W. and Covert, H. H. 2007. Diet of the Tonkin snub-nosed monkey (*Rhinopithecus avunculus*) in the Khau Ca area, Ha Giang Province, northeastern Vietnam. *Vietnamese Journal of Primatology* 1:75-83.
- [155] Li, Z. Y. and Rogers, M. E. 2006. Food items consumed by white-headed langurs in Fusui, China. *International Journal of Primatology* 27:1551-1567.
- [156] Matsuda, I., Tuuga, A. and Higashi, S. 2009. The feeding ecology and activity budget of proboscis monkeys. *American Journal of Primatology* 71:478-492.
- [157] Minhas, R. A., Ahmed, K. B., Awan, M. S. and Dar, N. I. 2010. Habitat utilization and feeding biology of Himalayan Grey Langur (*Semnopithecus entellus ajax*) in Machiara National Park, Azad Jammu and Kashmir, Pakistan. *Zoological Research* 2:177-188.
- [158] O'Brien, T. G. and Kinnaird, M. F. 1997. Behavior, diet, and movements of the Sulawesi crested black macaque (*Macaca nigra*). *International Journal of Primatology* 18:321-351.
- [159] Rawson, B. M. 2009. The socio-ecology of the black-shanked douc (*Pygathrix nigripes*) in Mondulkiri Province, Cambodia. *School of Archaeology and Anthropology*:211.
- [160] Su, H. H. and Lee, L. L. 2001. Food habits of Formosan rock macaques (*Macaca cyclopis*) in Jentse, northeastern Taiwan, assessed by fecal analysis and behavioral observation. *International Journal of Primatology* 22:359-377.
- [161] Workman, C. 2010. The foraging ecology of the Delacour's langur (*Trachypithecus delacouri*) in Van Long Nature Reserve, Vietnam. *Department of Evolutionary anthropology*:235.
- [162] Wu, J.-P., Zhou, W., Zhou, J.-L., Ai, H.-S., Huang, X.-X. and Li, J.-H. 2009. Diet and daily feeding amount of Hoolock Gibbon (*Hoolock hoolock*) at Nankang, Mt. Gaoligong. *Zoological Research* 30:539-544.
- [163] Zhang, X.-X., Zhou, W., Wu, J.-P., Bai, B., Li, Z.-B. and Li, J.-H. 2008. Food selection of Hoolock Gibbon (*Hoolock hoolock*) at Nankang, Mt. Gaoligong in spring. *Zoological Research* 29:174-180.
- [164] Zhou, Q., Wei, F., Li, M., Huang, C. and Luo, B. 2006. Diet and food choice of *Trachypithecus francoisi* in the Nonggang Nature Reserve, China. *International Journal of Primatology* 27:1441-1460.
- [165] Zhou, Q., Wei, H., Huang, Z. and Hunag, C. 2011. Diet of the Assamese macaque *Macaca assamensis* in limestone habitats of Nonggang, China. *Current Zoology* 57:18-25.

Appendix 1. Nutmeg-frugivore interactions in the Asia-Pacific region.

Plant species	Frugivore species	Frugivore group	Weight (kg)	Study site	Source	
<i>Endocomia macrocoma</i>	<i>Buceros bicornis</i>	Hornbill	2.5	Budo Sungapi Padi, Thailand	[57]	
	<i>Rhyticeros undulatus</i>	Hornbill	2.7	Budo Sungapi Padi, Thailand	[57]	
<i>Gymnacranthera farquhariana</i>	<i>R. undulatus</i>	Hornbill	2.7	Budo Sungapi Padi, Thailand	[57]	
	<i>Pongo pygmaeus</i>	Orangutan	57.5	Tuanan, Indonesia	[94]	
	<i>Hylobates klossii</i>	Gibbon	5.8	Siberut, Indonesia	[95]	
	<i>P. pygmaeus</i>	Orangutan	57.5	Suaq Balimbing, Indonesia	[94]	
	<i>Aceros cassidix</i>	Hornbill	2.4	Tangkoko DuaSudara, Indonesia	[63, 96]	
<i>Gymnacranthera</i> sp.GP	<i>P. pygmaeus</i>	Orangutan	57.5	Gunung Palu, Indonesia	[94]	
<i>Gymnacranthera</i> sp.LM001	<i>Anorrhinus galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]	
	<i>Anthracoseros malayanus</i>	Hornbill	1	Kutai, Indonesia	[45]	
<i>Gymnacranthera</i> sp.MKF001	<i>A. cassidix</i>	Hornbill	2.4	Tangkoko DuaSudara, Indonesia	[63, 96]	
<i>Horsfieldia amygdalina</i>	<i>Aceros nipalensis</i>	Hornbill	2.4	Huai Kha Khaeng, Thailand	[97]	
	<i>Anorrhinus austeni</i>	Hornbill	0.9	Khao Yai, Thailand	[10, 59, 98]	
	<i>Anthracoseros albirostris</i>	Hornbill	0.8	Khao Yai, Thailand	[10, 59, 98]	
	<i>B. bicornis</i>	Hornbill	2.5	Khao Yai, Thailand	[10, 59, 98]	
	<i>R. undulatus</i>	Hornbill	2.7	Khao Yai, Thailand	[10, 59, 98]	
	<i>Ducula badia</i>	Pigeon	0.6	Khao Yai, Thailand	[10]	
	<i>Callosciurus finlaysonii</i>	Squirrel	0.3	Khao Yai, Thailand	[10]	
	<i>A. galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]	
	<i>H. brachiata</i>	<i>Buceros rhinoceros</i>	Hornbill	2.8	Bukit Barisan Selatan, Indonesia	[99]
	<i>H. crassifolia</i>	<i>P. pygmaeus</i>	Orangutan	57.5	Sebangau, Indonesia	[100]
<i>Presbytis rubicunda</i>		Leaf monkey	6	Tanjung, Indonesia	[101]	
<i>P. pygmaeus</i>		Orangutan	57.5	Sebangau, Indonesia	[94]	
<i>P. pygmaeus</i>		Orangutan	57.5	Tuanan, Indonesia	[94]	
<i>H. glabra</i>	<i>P. pygmaeus</i>	Orangutan	57.5	Suaq Balimbing, Indonesia	[94]	
<i>H. grandis</i>	<i>P. pygmaeus</i>	Orangutan	57.5	Kubah, Malaysia	[94]	
<i>H. irya</i>	<i>Macaca fascicularis</i>	Macaque	4.5	Angaur, Micronesia	[102]	
	<i>Presbytis melalophos</i>	Leaf monkey	5.8	Krau, Malaysia	[103]	
	<i>M. fascicularis</i>	Macaque	4.5	Krau, Malaysia	[103]	
	Squirrel	Squirrel	NA	Krau, Malaysia	[103]	
	<i>Hylobates agilis</i>	Gibbon	6	Krau, Malaysia	[104]	
	<i>Ocyeros gingalensis</i>	Hornbill	0.2	Sinharaja, India	[105]	
	<i>Gallus lafayetti</i>	Junglefowl	0.6	Sinharaja, India	[105]	
	<i>Gracula ptilogenys</i>	Myna	0.2	Sinharaja, India	[105]	
<i>Moschiola meminna</i>	Mousedeer		7.5	Sinharaja, India	[105]	
	<i>Funambulus layardi</i>	Squirrel	0.1	Sinharaja, India	[105]	

	<i>B. bicornis</i>	Hornbill	2.5	Budo Sungapi Padi, Thailand	[57]
	<i>Rhyticeros plicatus</i>	Hornbill	1.6	Wide Bay, New Britain, PNG	[106]
<i>H. kingii</i>	<i>A. albirostris</i>	Hornbill	0.8	Pakke, India	[107]
	<i>B. bicornis</i>	Hornbill	2.5	Pakke, India	[107]
	<i>R. undulatus</i>	Hornbill	2.7	Pakke, India	[107]
	<i>D. badia</i>	Pigeon	0.6	Pakke, India	[108]
	<i>Trachypithecus shortridgei</i>	Leaf monkey	11.8	Pakke, India	[109]
<i>H. motleyi</i>	<i>Aceros comatus</i>	Hornbill	1.4	Kutai, Indonesia	[45]
	<i>A. galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]
	<i>B. rhinoceros</i>	Hornbill	2.8	Kutai, Indonesia	[45]
<i>H. pachyrachis</i>	<i>Macaca tonkeana</i>	Macaque	12	Lore Lindu, Indonesia	[110]
<i>H. reticulata</i>	<i>A. galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]
<i>H. sucosa</i>	<i>Callosciurus</i> sp.	Squirrel	NA	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)
	<i>P. melalophos</i>	Leaf monkey	5.8	Krau, Malaysia	[103]
	Squirrel	Squirrel	NA	Krau, Malaysia	[103]
	<i>A. comatus</i>	Hornbill	1.4	Budo Sungapi Padi, Thailand	[57]
	<i>Aceros corrugates</i>	Hornbill	1.6	Budo Sungapi Padi, Thailand	[57]
	<i>B. bicornis</i>	Hornbill	2.5	Budo Sungapi Padi, Thailand	[57]
	<i>R. undulatus</i>	Hornbill	2.7	Budo Sungapi Padi, Thailand	[57]
	<i>Presbytis femoralis</i>	Leaf monkey	7.3	Pasoh, Malaysia	[58]
	<i>Hystrix brachyura</i>	Porcupine	8	Pasoh, Malaysia	[58]
<i>H. superba</i>	Squirrel	Squirrel	NA	Krau, Malaysia	[103]
	<i>Macaca nemestrina</i>	Macaque	8.9	Pasoh, Malaysia	[58]
	<i>Leopoldamys sabanus</i>	Rodent	0.4	Pasoh, Malaysia	[58]
	<i>Maxomys surifer</i>	Rodent	0.2	Pasoh, Malaysia	[58]
	<i>Lariscus insignis</i>	Squirrel	0.2	Pasoh, Malaysia	[58]
	<i>Varanus bengalensis</i>	Lizard	7	Pasoh, Malaysia	[58]
<i>H. tomentosa</i>	<i>Trachypithecus obscurus</i>	Leaf monkey	7	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)
	<i>Callosciurus</i> sp.	Squirrel	NA	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)
	<i>A. corrugates</i>	Hornbill	1.6	Budo Sungapi Padi, Thailand	[57]
	<i>A. galeritus</i>	Hornbill	1.1	Budo Sungapi Padi, Thailand	[57]
	<i>B. bicornis</i>	Hornbill	2.5	Budo Sungapi Padi, Thailand	[57]
	<i>Buceros vijil</i>	Hornbill	2.9	Budo Sungapi Padi, Thailand	[57]
	<i>R. undulatus</i>	Hornbill	2.7	Budo Sungapi Padi, Thailand	[57]
<i>H. wallichii</i>	<i>A. galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]
<i>Horsfieldia</i> sp.CH051	<i>P. melalophos</i>	Leaf monkey	5.8	Krau, Malaysia	[103]

	<i>M. fascicularis</i>	Macaque	4.5	Krau, Malaysia	[103]
	Squirrel	Squirrel	NA	Krau, Malaysia	[103]
<i>Horsfieldia</i> sp.FR001	<i>Helarctos malayanus</i>	Bear	46	Sungai Wain, Indonesia	[111]
<i>Knema attenuata</i>	<i>Ocyrceros griseus</i>	Hornbill	0.3	Indira Gandhi, India	[112]
<i>K. attenuata</i>	<i>Paradoxurus jerdoni</i>	Civet	4	Kalakad Mundauthurai, India	[113]
	<i>Semnopithecus entellus</i>	Leaf monkey	13.8	Maharashtra, India	[114]
	<i>Macaca silenus</i>	Macaque	7.5	Indira Gandhi, India	[115]
<i>K. cinerea</i>	<i>A. albirostris</i>	Hornbill	0.8	Pakke, India	[108]
	<i>B. bicornis</i>	Hornbill	2.5	Pakke, India	[108]
	<i>R. undulatus</i>	Hornbill	2.7	Pakke, India	[108]
	<i>D. badia</i>	Pigeon	0.6	Pakke, India	[108]
	<i>Hylobates lar</i>	Gibbon	6	Krau, Malaysia	[103]
	<i>Symphalangus syndactylus</i>	Gibbon	12.4	Krau, Malaysia	[103]
	Squirrel	Squirrel	NA	Krau, Malaysia	[103]
	<i>Hylobates moloch</i>	Gibbon	5.7	Gunung Halimun, Indonesia	[116]
	<i>Presbytis comata</i>	Leaf monkey	6.5	Gunung Halimun, Indonesia	[116]
	<i>Trachypithecus auratus</i>	Leaf monkey	7.1	Gunung Halimun, Indonesia	[116]
	<i>M. tonkeana</i>	Macaque	12	Lore Lindu, Indonesia	[110]
	<i>P. pygmaeus</i>	Orangutan	57.5	Ketambe, Indonesia	[94]
	<i>P. pygmaeus</i>	Orangutan	57.5	Ketambe, Indonesia	[94]
<i>K. conferta</i>	<i>H. klossii</i>	Gibbon	5.8	Siberut, Indonesia	[95]
<i>K. curtisii</i>	<i>Callosciurus</i> sp.	Squirrel	NA	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)
	<i>A. galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]
<i>K. elegans</i>	<i>H. lar</i>	Gibbon	6	Khao Yai, Thailand	[117]
	<i>A. austeni</i>	Hornbill	0.9	Khao Yai, Thailand	[10, 59, 98]
	<i>A. albirostris</i>	Hornbill	0.8	Khao Yai, Thailand	[10, 59, 98]
	<i>B. bicornis</i>	Hornbill	2.5	Khao Yai, Thailand	[10, 59, 98]
	<i>R. undulatus</i>	Hornbill	2.7	Khao Yai, Thailand	[10, 59, 98]
	<i>D. badia</i>	Pigeon	0.6	Khao Yai, Thailand	[10]
	<i>C. finlaysonii</i>	Squirrel	0.3	Khao Yai, Thailand	[10]
	<i>Hylobates pileatus</i>	Gibbon	8.4	Khao Yai, Thailand	[118]
	<i>Paradoxurus hermaphroditus</i>	Civet	3.2	Khao Yai, Thailand	[51]
	<i>Muntiacus muntjak</i>	Deer	24	Khao Yai, Thailand	[51]
	<i>M. surifer</i>	Rodent	0.2	Khao Yai, Thailand	[51]
	<i>C. finlaysonii</i>	Squirrel	0.3	Khao Yai, Thailand	[51]
	<i>Viverra zibetha</i>	Civet	6.3	Khao Yai, Thailand	[71]
	<i>H. lar</i>	Gibbon	6	Khao Yai, Thailand	[71, 119]
	<i>Atherurus macrourus</i>	Porcupine	6	Khao Yai, Thailand	[71, 119]

<i>K. erratica</i>	<i>A. nipalensis</i>	Hornbill	2.4	Huai Kha Khaeng, Thailand	[97]	
<i>K. furfuracea</i>	<i>T. obscurus</i>	Leaf monkey	7	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)	
	<i>Callosciurus</i> sp.	Squirrel	NA	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)	
	<i>A. comatus</i>	Hornbill	1.4	Budo Sungapi Padi, Thailand	[57]	
	<i>B. bicornis</i>	Hornbill	2.5	Budo Sungapi Padi, Thailand	[57]	
	<i>B. rhinoceros</i>	Hornbill	2.8	Budo Sungapi Padi, Thailand	[57]	
	<i>R. undulatus</i>	Hornbill	2.7	Budo Sungapi Padi, Thailand	[57]	
	<i>K. glauca</i>	<i>Callosciurus</i> sp.	Squirrel	NA	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)
<i>K. glaucescens</i>	<i>P. pygmaeus</i>	Orangutan	57.5	Sungai Wain, Indonesia	[94]	
<i>K. globularia</i>	<i>T. obscurus</i>	Leaf monkey	7	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)	
	<i>Callosciurus</i> sp.	Squirrel	NA	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)	
	<i>A. corrugates</i>	Hornbill	1.6	Budo Sungapi Padi, Thailand	[57]	
	<i>A. galeritus</i>	Hornbill	1.1	Budo Sungapi Padi, Thailand	[57]	
	<i>B. bicornis</i>	Hornbill	2.5	Budo Sungapi Padi, Thailand	[57]	
	<i>R. undulatus</i>	Hornbill	2.7	Budo Sungapi Padi, Thailand	[57]	
<i>K. hookeriana</i>	<i>Callosciurus</i> sp.	Squirrel	NA	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)	
	<i>H. lar</i>	Gibbon	6	Krau, Malaysia	[103]	
	<i>S. syndactylus</i>	Gibbon	12.4	Krau, Malaysia	[103]	
	Squirrel	Squirrel	NA	Krau, Malaysia	[103]	
	<i>A. galeritus</i>	Hornbill	1.1	Budo Sungapi Padi, Thailand	[57]	
	<i>M. nemestrina</i>	Macaque	8.9	Pasoh, Malaysia	[58]	
	<i>Sus scrofa</i>	Pig	70	Pasoh, Malaysia	[58]	
	<i>Trichys fasciculata</i>	Porcupine	2.5	Pasoh, Malaysia	[58]	
	<i>L. sabanus</i>	Rodent	0.4	Pasoh, Malaysia	[58]	
	<i>M. surifer</i>	Rodent	0.2	Pasoh, Malaysia	[58]	
	<i>L. insignis</i>	Squirrel	0.2	Pasoh, Malaysia	[58]	
	<i>K. intermedia</i>	<i>P. rubicunda</i>	Leaf monkey	6	Tanjung, Indonesia	[101]
	<i>K. latericia</i>	<i>H. malayanus</i>	Bear	46	Sungai Wain, Indonesia	[111]
<i>P. pygmaeus</i>		Orangutan	57.5	Gunung Palu, Indonesia	[120]	
<i>A. cassidix</i>		Hornbill	2.4	Tangkoko DuaSudara, Indonesia	[63, 96]	
<i>A. galeritus</i>		Hornbill	1.1	Kutai, Indonesia	[45]	
<i>A. galeritus</i>		Hornbill	1.1	Kutai, Indonesia	[45]	
<i>R. undulatus</i>		Hornbill	2.7	Kutai, Indonesia	[45]	

	<i>M. fascicularis</i>	Macaque	4.5	Bukit Timah, Singapore	[121]
	<i>P. pygmaeus</i>	Orangutan	57.5	Mentoko, Indonesia	[94]
	<i>P. pygmaeus</i>	Orangutan	57.5	Meratus, Indonesia	[94]
<i>K. latifolia</i>	<i>Hylobates muellerixagilis</i>	Gibbon	5.7	Barito Ulu, Indonesia	[122]
<i>K. laurina</i>	<i>P. pygmaeus</i>	Orangutan	57.5	Danum Valley, Malaysia	[123]
	<i>M. fascicularis</i>	Macaque	4.5	Bukit Timah, Singapore	[121]
	<i>A. nipalensis</i>	Hornbill	2.4	Huai Kha Khaeng, Thailand	[124, 125]
	<i>B. bicornis</i>	Hornbill	2.5	Huai Kha Khaeng, Thailand	[124, 125]
	<i>Presbytis thomasi</i>	Leaf monkey	6.7	Ketambe, Indonesia	[126]
	<i>P. pygmaeus</i>	Orangutan	57.5	Ketambe, Indonesia	[94]
<i>K. pallens</i>	<i>A. galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]
<i>K. percoriacea</i>	<i>H. muellerixagilis</i>	Gibbon	5.7	Barito Ulu, Indonesia	[122]
<i>K. pseudolaurina</i>	<i>T. obscurus</i>	Leaf monkey	7	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)
	<i>Callosciurus</i> sp.	Squirrel	NA	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)
	<i>B. bicornis</i>	Hornbill	2.5	Budo Sungapi Padi, Thailand	[57]
<i>K. scortechinii</i>	<i>M. nemestrina</i>	Macaque	8.9	Pasoh, Malaysia	[58]
	<i>Tragulus kanchil</i>	Mousedeer	2.3	Pasoh, Malaysia	[58]
	<i>H. brachyura</i>	Porcupine	8	Pasoh, Malaysia	[58]
	<i>Rhinosciurus laticaudatus</i>	Squirrel	0.2	Pasoh, Malaysia	[58]
	<i>Tupaia glis</i>	Tupaia	0.2	Pasoh, Malaysia	[58]
<i>Knema</i> sp.CH052	<i>Calyptomena viridis</i>	Broadbill	0.1	Krau, Malaysia	[103]
	Hornbill 6 spp.	Hornbill	NA	Krau, Malaysia	[103]
	<i>P. melalophos</i>	Leaf monkey	5.8	Krau, Malaysia	[103]
	<i>Callosciurus prevostii</i>	Squirrel	0.4	Krau, Malaysia	[103]
<i>Knema</i> sp.FR002	<i>H. malayanus</i>	Bear	46	Sungai Wain, Indonesia	[111]
<i>Knema</i> sp.GI001	<i>H. agilis</i>	Gibbon	6	Krau, Malaysia	[104]
<i>Knema</i> sp.GP	<i>P. pygmaeus</i>	Orangutan	57.5	Gunung Palu, Indonesia	[94]
<i>Knema</i> sp.KA001	<i>P. pygmaeus</i>	Orangutan	57.5	Danum Valley, Malaysia	[123]
<i>Knema</i> sp.KO001	<i>A. nipalensis</i>	Hornbill	2.4	Huai Kha Khaeng, Thailand	[124, 125]
<i>Knema</i> sp.ME	<i>P. pygmaeus</i>	Orangutan	57.5	Meratus, Indonesia	[94]
<i>Knema</i> sp.SW	<i>P. pygmaeus</i>	Orangutan	57.5	Sungai Wain, Indonesia	[94]
<i>Knema</i> sp.US	<i>P. pygmaeus</i>	Orangutan	57.5	Ulu Segama, Malaysia	[94]
<i>Knema</i> sp.VC001	<i>A. nipalensis</i>	Hornbill	2.4	Huai Kha Khaeng, Thailand	[97]
	<i>Rhyticeros subruficollis</i>	Hornbill	2.1	Huai Kha Khaeng, Thailand	[97]
<i>Knema</i> sp.YH001	<i>A. galeritus</i>	Hornbill	1.1	Bukit Barisan Selatan, Indonesia	[99]
<i>Knema</i> sp.YH002	<i>B. rhinoceros</i>	Hornbill	2.8	Bukit Barisan Selatan, Indonesia	[99]
	<i>R. undulatus</i>	Hornbill	2.7	Bukit Barisan Selatan, Indonesia	[99]

<i>Myristica andamanica</i>	<i>Ducula aenea</i>	Pigeon	0.5	NA	[127]	
<i>M. beddomii</i>	<i>M. silenus</i>	Macaque	7.5	Indira Gandhi, India	[115]	
<i>M. ceylanica</i>	<i>Aceros waldeni</i>	Hornbill	1.1	North Negros, Philippines	[11]	
	<i>Penelopides panini</i>	Hornbill	0.5	North Negros, Philippines	[11]	
	<i>A. waldeni</i>	Hornbill	1.1	North Negros, Philippines	[128]	
<i>M. cinnamomea</i>	<i>P. pygmaeus</i>	Orangutan	57.5	Ulu Segama, Malaysia	[94]	
	<i>M. nemestrina</i>	Macaque	8.9	Pasoh, Malaysia	[58]	
	<i>H. brachyura</i>	Porcupine	8	Pasoh, Malaysia	[58]	
	<i>T. fasciculata</i>	Porcupine	2.5	Pasoh, Malaysia	[58]	
	<i>M. surifer</i>	Rodent	0.2	Pasoh, Malaysia	[58]	
	<i>R. laticaudatus</i>	Squirrel	0.2	Pasoh, Malaysia	[58]	
<i>M. dactyloides</i>	<i>V. bengalensis</i>	Lizard	7	Pasoh, Malaysia	[58]	
	<i>Semnopithecus johnii</i>	Leaf monkey	11	Kalakad Mundauthurai, India	[129]	
	<i>M. silenus</i>	Macaque	7.5	Kalakad Mundauthurai, India	[129]	
	<i>Platacanthomys lasiurus</i>	Rodent	0.8	Kalakad Mundauthurai, India	[129]	
	<i>Ratufa indica</i>	Squirrel	2	Kalakad Mundauthurai, India	[129]	
	<i>O. gingalensis</i>	Hornbill	0.2	Sinharaja, India	[105]	
	<i>G. ptilogenys</i>	Myna	0.2	Sinharaja, India	[105]	
	<i>Viverricula indica</i>	Civet	3	Sinharaja, India	[105]	
	<i>Herpestes fuscus</i>	Mongoose	2.7	Sinharaja, India	[105]	
	<i>F. layardi</i>	Squirrel	0.1	Sinharaja, India	[105]	
	<i>O. griseus</i>	Hornbill	0.3	Indira Gandhi, India	[112]	
	<i>M. silenus</i>	Macaque	7.5	Indira Gandhi, India	[130]	
	<i>M. elliptica</i>	<i>A. comatus</i>	Hornbill	1.4	Budo Sungapi Padi, Thailand	[57]
		<i>A. galeritus</i>	Hornbill	1.1	Budo Sungapi Padi, Thailand	[57]
		<i>R. undulatus</i>	Hornbill	2.7	Budo Sungapi Padi, Thailand	[57]
<i>M. nemestrina</i>		Macaque	8.9	Pasoh, Malaysia	[58]	
<i>H. brachyura</i>		Porcupine	8	Pasoh, Malaysia	[58]	
<i>T. fasciculata</i>		Porcupine	2.5	Pasoh, Malaysia	[58]	
<i>L. sabanus</i>		Rodent	0.4	Pasoh, Malaysia	[58]	
<i>M. surifer</i>		Rodent	0.2	Pasoh, Malaysia	[58]	
<i>Maxomys whiteheadi</i>		Rodent	0.6	Pasoh, Malaysia	[58]	
<i>R. laticaudatus</i>		Squirrel	0.2	Pasoh, Malaysia	[58]	
<i>M. fatua</i>	<i>V. bengalensis</i>	Lizard	7	Pasoh, Malaysia	[58]	
	<i>R. plicatus</i>	Hornbill	1.6	Crater Mountain, PNG	[131]	
<i>M. guatteriifolia</i>	<i>Rhyticeros everetti</i>	Hornbill	1.1	Sumba, Indonesia	[132]	
<i>M. hypargyrea</i>	<i>Ducula pacifica</i>	Pigeon	0.4	Eua, Tonga	[70]	
	<i>Pteropus tonganus</i>	Flying fox	0.6	Eua, Tonga	[62]	
	<i>Prosopieia tabuensis</i>	Parrot	0.2	Eua, Tonga	[62]	

	<i>Rattus</i> spp.	Rodent	0.2	Eua, Tonga	[62, 133]
<i>M. iners</i>	<i>Callosciurus</i> sp.	Squirrel	NA	Budo Sungapi Padi, Thailand	(S. Kitamura, unpublished data)
	<i>A. comatus</i>	Hornbill	1.4	Budo Sungapi Padi, Thailand	[57]
	<i>A. galeritus</i>	Hornbill	1.1	Budo Sungapi Padi, Thailand	[57]
	<i>B. bicornis</i>	Hornbill	2.5	Budo Sungapi Padi, Thailand	[57]
	<i>B. rhinoceros</i>	Hornbill	2.8	Budo Sungapi Padi, Thailand	[57]
	<i>R. undulatus</i>	Hornbill	2.7	Budo Sungapi Padi, Thailand	[57]
<i>M. lowiana</i>	<i>P. pygmaeus</i>	Orangutan	57.5	Sebangau, Indonesia	[100]
	<i>P. pygmaeus</i>	Orangutan	57.5	Tuanan, Indonesia	[94]
<i>M. maingayi</i>	Squirrel	Squirrel	NA	Krau, Malaysia	[103]
	<i>M. nemestrina</i>	Macaque	8.9	Pasoh, Malaysia	[58]
	<i>T. kanchil</i>	Mousedeer	2.3	Pasoh, Malaysia	[58]
	<i>T. fasciculata</i>	Porcupine	2.5	Pasoh, Malaysia	[58]
	<i>L. sabanus</i>	Rodent	0.4	Pasoh, Malaysia	[58]
	<i>T. glis</i>	Tupai	0.2	Pasoh, Malaysia	[58]
<i>M. malabarica</i>	<i>S. entellus</i>	Leaf monkey	14.5	Maharashtra, India	[114]
	<i>M. silenus</i>	Macaque	7.5	Indira Gandhi, India	[130]
<i>M. maxima</i>	<i>R. undulatus</i>	Hornbill	2.7	Bukit Barisan Selatan, Indonesia	[99]
	<i>P. pygmaeus</i>	Orangutan	57.5	Sungai Wain, Indonesia	[94]
<i>M. muelleri</i>	<i>Ptilinopus magnificus</i>	Pigeon	0.2	Laceys Creek, Australia	[134]
	<i>Ptilinopus superbus</i>	Pigeon	0.1	Laceys Creek, Australia	[134]
	<i>Ducula spilorrhoa</i>	Pigeon	0.5	Laceys Creek, Australia	[134]
<i>M. subalulata</i>	<i>Manucodia chalybatus</i>	Bird of Paradise	0.2	Crater Mountain, PNG	[131]
	<i>Parotia lawesii</i>	Bird of Paradise	0.2	Crater Mountain, PNG	[131]
<i>M. venisa</i>	<i>H. muellerixagilis</i>	Gibbon	5.7	Barito Ulu, Indonesia	[122]
<i>M. villosa</i>	<i>P. pygmaeus</i>	Orangutan	57.5	Kinabatangan, Malaysia	[94]
	<i>P. pygmaeus</i>	Orangutan	57.5	Sungai Wain, Indonesia	[94]
<i>M. warburgii</i>	<i>R. plicatus</i>	Hornbill	1.6	Wide Bay, New Britain, PNG	[106]
<i>Myristica</i> sp.AL001	<i>D. aenea</i>	Pigeon	0.5	NA	[127]
<i>Myristica</i> sp.AL003	<i>Ducula bicolor</i>	Pigeon	0.5	NA	[127]
<i>Myristica</i> sp.AL004	<i>D. badia</i>	Pigeon	0.6	NA	[127]
	<i>A. nipalensis</i>	Hornbill	2.4	NA	[135]
<i>Myristica</i> sp.AL005	<i>B. bicornis</i>	Hornbill	2.5	NA	[135]
<i>Myristica</i> sp.BB001	<i>Diphyllodes magnificus</i>	Bird of Paradise	0.2	Mt. Missim, PNG	[136]
	<i>Manucodia keraudrenii</i>	Bird of Paradise	0.2	Mt. Missim, PNG	[136]
	<i>Paradisaea raggiana</i>	Bird of Paradise	0.2	Mt. Missim, PNG	[136]

	<i>P. lawesii</i>	Bird of Paradise	0.2	Mt. Missim, PNG	[136]
<i>Myristica</i> sp.BB002	<i>Cicinnurus magnificus</i>	Bird of Paradise	0.1	Varirata, PNG	[21]
	<i>M. chalybatus</i>	Bird of Paradise	0.2	Varirata, PNG	[21]
	<i>P. raggiana</i>	Bird of Paradise	0.2	Varirata, PNG	[21]
	<i>P. magnificus</i>	Pigeon	0.2	Varirata, PNG	[21]
	<i>Ptilinopus pulchellus</i>	Pigeon	0.1	Varirata, PNG	[21]
	<i>P. superbus</i>	Pigeon	0.1	Varirata, PNG	[21]
<i>Myristica</i> sp.CH053	<i>Hornbill 6spp</i>	Hornbill	NA	Krau, Malaysia	[103]
	<i>P. melalophos</i>	Leaf monkey	5.8	Krau, Malaysia	[103]
	<i>C. prevostii</i>	Squirrel	0.4	Krau, Malaysia	[103]
<i>Myristica</i> sp.CH054	Squirrel	Squirrel	NA	Krau, Malaysia	[103]
<i>Myristica</i> sp.GI003	<i>D. pacifica</i>	Pigeon	0.4	NA	[37]
<i>Myristica</i> sp.GI004	<i>Ducula latrans</i>	Pigeon	0.6	NA	[37]
<i>Myristica</i> sp.KET	<i>P. pygmaeus</i>	Orangutan	57.5	Ketambe, Indonesia	[94]
<i>Myristica</i> sp.KN	<i>P. pygmaeus</i>	Orangutan	57.5	Kinabatangan, Malaysia	[94]
<i>Myristica</i> sp.LM001	<i>A. galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]
<i>Myristica</i> sp.LM002	<i>A. comatus</i>	Hornbill	1.4	Kutai, Indonesia	[45]
	<i>A. corrugates</i>	Hornbill	1.6	Kutai, Indonesia	[45]
	<i>A. galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]
	<i>B. rhinoceros</i>	Hornbill	2.8	Kutai, Indonesia	[45]
<i>Myristica</i> sp.LM003	<i>A. comatus</i>	Hornbill	1.4	Kutai, Indonesia	[45]
	<i>B. rhinoceros</i>	Hornbill	2.8	Kutai, Indonesia	[45]
	<i>R. undulatus</i>	Hornbill	2.7	Kutai, Indonesia	[45]
<i>Myristica</i> sp.LM004	<i>A. comatus</i>	Hornbill	1.4	Kutai, Indonesia	[45]
	<i>A. galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]
	<i>R. undulatus</i>	Hornbill	2.7	Kutai, Indonesia	[45]
<i>Myristica</i> sp.LM005	<i>A. comatus</i>	Hornbill	1.4	Kutai, Indonesia	[45]
	<i>A. galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]
	<i>A. malayanus</i>	Hornbill	1	Kutai, Indonesia	[45]
	<i>R. undulatus</i>	Hornbill	2.7	Kutai, Indonesia	[45]
<i>Myristica</i> sp.ML006	<i>A. comatus</i>	Hornbill	1.4	Kutai, Indonesia	[45]
	<i>A. corrugates</i>	Hornbill	1.6	Kutai, Indonesia	[45]
	<i>A. galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]
	<i>A. malayanus</i>	Hornbill	1	Kutai, Indonesia	[45]
	<i>B. rhinoceros</i>	Hornbill	2.8	Kutai, Indonesia	[45]
<i>Myristica</i> sp.ML007	<i>A. comatus</i>	Hornbill	1.4	Kutai, Indonesia	[45]
	<i>A. galeritus</i>	Hornbill	1.1	Kutai, Indonesia	[45]
	<i>B. rhinoceros</i>	Hornbill	2.8	Kutai, Indonesia	[45]

	<i>R. undulatus</i>	Hornbill	2.7	Kutai, Indonesia	[45]
<i>Myristica</i> sp.PA001	<i>H. lar</i>	Gibbon	6	Ketambe, Indonesia	[137]
<i>Myristica</i> sp.RK001	<i>B. bicornis</i>	Hornbill	2.5	Indira Gandhi, India	[46]
	<i>O. griseus</i>	Hornbill	0.3	Indira Gandhi, India	[46]
	<i>D. badia</i>	Pigeon	0.6	Indira Gandhi, India	[46]
<i>Myristica</i> sp.SM001	<i>D. bicolor</i>	Pigeon	0.5	NA	[138]
<i>Myristica</i> sp.TN	<i>P. pygmaeus</i>	Orangutan	57.5	Tuanan, Indonesia	[94]
<i>Myristica</i> sp.US	<i>P. pygmaeus</i>	Orangutan	57.5	Ulu Segama, Malaysia	[94]
Myristicaceae sp.AZ001	<i>Trachypithecus phayrei</i>	Leaf monkey	7.4	Lawachara, Bangladesh	[139]
Myristicaceae sp.BL001	<i>Nomascus concolor</i>	Gibbon	6.8	Wu Liang, China	[140]
Myristicaceae sp.DE001	<i>Trachypithecus vetulus</i>	Leaf monkey	8.2	Panadura, Sri Lanka	[141]
Myristicaceae sp.DU001	<i>Pygathrix nigripes</i>	Leaf monkey	10.7	Phuoc Binh, Vietnam	[142]
Myristicaceae sp.FA001	<i>N. concolor</i>	Gibbon	6.8	Dazhaizi, China	[143]
Myristicaceae sp.FR001	<i>Ptilinopus iozonus</i>	Pigeon	0.1	Brown River, PNG	[144]
	<i>P. magnificus</i>	Pigeon	0.2	Brown River, PNG	[144]
	<i>P. superbus</i>	Pigeon	0.1	Brown River, PNG	[144]
	<i>P. pulchellus</i>	Pigeon	0.1	Brown River, PNG	[144]
	<i>Ducula rufigaster</i>	Pigeon	0.5	Brown River, PNG	[144]
Myristicaceae sp.FU001	<i>Presbytis potenziani</i>	Leaf monkey	6.4	Betumonga, Indonesia	[145]
Myristicaceae sp.GI002	<i>D. aenea</i>	Pigeon	0.5	NA	[37]
Myristicaceae sp.GO001	<i>D. aenea</i>	Pigeon	0.5	NA	[146]
Myristicaceae sp.GO002	<i>D. pacifica</i>	Pigeon	0.4	NA	[146]
Myristicaceae sp.GO003	<i>D. bicolor</i>	Pigeon	0.5	NA	[146]
Myristicaceae sp.GO004	<i>D. spilorrhoea</i>	Pigeon	0.5	NA	[146]
Myristicaceae sp.GR001	<i>Macaca mulatta</i>	Macaque	9.9	Baimaxueshan, China	[147]
Myristicaceae sp.GU001	<i>P. thomasi</i>	Leaf monkey	6.7	Gunung Leuser, Indonesia	[148]
Myristicaceae sp.IS001	<i>Hoolock hoolock</i>	Gibbon	6.5	Lawachara & Chunut, Bangladesh	[149]
Myristicaceae sp.KET	<i>P. pygmaeus</i>	Orangutan	57.5	Ketambe, Indonesia	[94]
Myristicaceae sp.KO001	<i>T. auratus</i>	Leaf monkey	7.1	Pangandaran, Indonesia	[150]
Myristicaceae sp.KR001	<i>Macaca radiata</i>	Macaque	5.3	Marakkanam, India	[151]
Myristicaceae sp.KU001	<i>Trachypithecus pileatus</i>	Leaf monkey	11.4	Pakke, India	[152]
Myristicaceae sp.KU002	<i>Macaca munzala</i>	Macaque	15	Zemithang, India	[153]
Myristicaceae sp.LE001	<i>Rhinopithecus avunculus</i>	Leaf monkey	11.3	Khau Ca, Vietnam	[154]
Myristicaceae sp.LI001	<i>Trachypithecus poliocephalus</i>	Leaf monkey	8.1	Fusui, China	[155]
Myristicaceae sp.MA001	<i>Nasalis larvatus</i>	Leaf monkey	15.6	Kinabatangan, Malaysia	[156]
Myristicaceae sp.MI001	<i>Semnopithecus ajax</i>	Leaf monkey	14.9	Machiara, Pakistan	[157]
Myristicaceae sp.OB001	<i>Macaca nigra</i>	Macaque	7.7	Tangkoko DuaSudara, Indonesia	[158]
Myristicaceae sp.RA001	<i>P. nigripes</i>	Leaf monkey	10.7	Seima, Cambodia	[159]
Myristicaceae sp.SM001	<i>R. plicatus</i>	Hornbill	1.6	Wide Bay, New Britain, PNG	[106]

Myristicaceae sp.SU001	<i>Macaca cyclopis</i>	Macaque	5.5	Jentse, Taiwan	[160]
Myristicaceae sp.TM001	<i>Macaca arctoides</i>	Macaque	10.3	Khao Krapuk, Thailand	(T. Maruhashi, unpublished data)
Myristicaceae sp.WO001	<i>Trachypithecus delacouri</i>	Leaf monkey	8.4	Van Long, Vietnam	[161]
Myristicaceae sp.WU001	<i>H. hoolock</i>	Gibbon	6.5	Nankang, China	[162, 163]
Myristicaceae sp.ZH001	<i>Trachypithecus francoisi</i>	Leaf monkey	5.9	Nonggang, China	[164]
Myristicaceae sp.ZH002	<i>Macaca assamensis</i>	Macaque	9.1	Nonggang, China	[165]